

Baryon-Baryon Bound States From First Principles

Paulo A. Faria da Veiga* and Michael O'Carroll

Departamento de Matemática Aplicada e Estatística, ICMC-USP

C.P. 668, 13560-970 São Carlos SP, Brazil.

(Dated: October 10, 2006)

We determine baryon-baryon bound states in $3 + 1$ dimensional $SU(3)$ lattice QCD with two flavors, 4×4 spin matrices, and in an imaginary time formulation. For small hopping parameter, $\kappa > 0$, and large glueball mass (strong coupling), we show the existence of three-quark isospin $1/2$ particles (proton and neutron) and isospin $3/2$ baryons (delta particles), with asymptotic masses $-3 \ln \kappa$ and isolated dispersion curves. We only consider the existence of bound states of total isospin $I = 0, 3$. Using a ladder approximation to a lattice Bethe-Salpeter equation, baryon-baryon bound states are found in these two sectors, with asymptotic masses $-6 \ln \kappa$ and binding energies of order κ^2 . The dominant baryon-baryon interaction is an energy-independent spatial range-one potential with an $\mathcal{O}(\kappa^2)$ strength. There is also attraction arising from gauge field correlations associated with six overlapping bonds, but it is counterbalanced by Pauli repulsion to give a vanishing zero-range potential. The overall range-one potential results from a quark, antiquark exchange with *no* meson exchange interpretation; the repulsive or attractive nature of the interaction does depend on the isospin and spin of the two-baryon state.

PACS numbers: 11.15.Ha, 02.30.Tb, 11.10.St, 24.85.+p.

Keywords: Excitation Spectrum, Lattice QCD, Bethe-Salpeter Equation, Spectral Analysis

1. INTRODUCTION AND RESULTS

Our aim in this paper is to obtain the low-lying energy-momentum (EM) spectrum such as particles and their bound states in $SU(3)$ lattice quantum chromodynamics (QCD) [1, 2, 3, 13]. We treat a lattice QCD model in $3 + 1$ dimensions, with 4×4 Dirac spin matrices, and with two-flavor quarks (*up* and *down*). This model has a global $SU(2)$ flavor (isospin) symmetry, and accommodates twenty, three-quark, one-particle states (baryons), and their antiparticles, associated with the proton (*p*), the neutron (*n*) and the delta (Δ) particles.

This work is an important part of a recent series of papers given in Refs. [4, 5, 6, 7, 8, 9, 10] where, treating less complex QCD models, we started a research program aiming at developing a building up process and techniques to understand, from first principles, the hadronic particles and their bound states, bridging the gap between QCD and nuclear physics. The existence of baryons and mesons is a manifestation of confinement. Our main goal is to understand the mechanisms responsible for attraction and repulsion between the hadronic particles, when and how bound states occur and how their binding is related to the effective Yukawa meson-exchange theory.

Here, as in previous papers (see Refs. [4, 5, 6, 7, 8, 9, 10]), the analysis is performed in an imaginary-time formulation of lattice QCD, and is restricted to the strong coupling regime (small hopping parameter $0 < \kappa \ll 1$, and large glueball mass $\beta \equiv 1/g_0^2 \ll \kappa$). Even though the strong coupling regime is far from the scaling limit

(continuum quantum field theory), these models do exhibit particles and we hope, as for confinement, that the main features of attraction and repulsion of the continuum model are present in the strong coupling regime.

Of course, there is the pointwise repulsive interaction originating from Pauli exclusion. As expected, there is also a nonlocal interaction arising from a quark-antiquark (*not* a meson particle!) exchange mechanism like in the effective Yukawa theory. However, the attractive or repulsive nature of this interaction does depend on the spin and isospin of the two-baryon state. How this dependence occurs is to be understood. Not less important, our analysis displays another essential mechanism of attraction arising from gauge field correlation effects associated with integration over (six) overlapping gauge bonds: if these gauge correlations are not present the bound states cease to exist.

Our general approach to detect particles and bound states in the low-lying part of the spectrum is to derive spectral representations for the two- and four-baryon functions which allow us to relate their complex momentum singularities to the EM spectrum via a Feynman-Kac (F-K) formula. For other approaches to the determination of the spectrum both at the theoretical and numerical levels, see Refs. [1, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21] and Refs. [22, 23, 24, 25] dealing with numerical simulations.

It must be pointed out that the detection of particle (and their bound states) masses through the exponential decay rate of suitable correlations, *without* a spectral representation, is meaningless, and the resulting values may be far from the correct ones, especially in cases where degeneracies are broken with small spectral separations.

To determine the bound state spectrum, in a first step, we use a ladder approximation to a lattice version of the Bethe-Salpeter (B-S) equation for suitable four-baryon

*Electronic address: veiga@icmc.usp.br

correlations. However, our work does incorporate the main ingredients to go beyond this ladder approximation and analytically validate our results to the full model in the strong coupling regime, similar to what is done in some classical and stochastic spin systems (see Refs. [26, 27]).

In our previous papers, the simplest case in which a two-baryon bound state appeared is in the two-flavor total isospin $I = 0, 1$ sectors with 2×2 spin matrices in $2+1$ dimensions (see Ref. [9]); there are *no* bound states for $I = 2, 3$. However, this model is not complex enough to accommodate states like protons and neutrons in the one-particle spectrum.

Here, considering a more realistic lattice QCD model, it is shown that there are twenty one-baryon states with asymptotic masses given by $-3 \ln \kappa - 3\kappa^3/4 + \mathcal{O}(\kappa^6)$ and isolated dispersion curves (the upper gap property). The upper gap property is unknown in the Hamiltonian formulation of the model.

Concerning the two-particle spectrum, we determine the two-baryon bound states for the total isospin $I = 0$ and $I = 3$ sectors and below the two-baryon threshold, which is given by twice the smallest of the baryon mass. We show that there are diverse two-baryon bound states. Their asymptotic masses are $-6 \ln \kappa$ and their binding energies are of order κ^2 . In the $I = 0$ sector, the most strongly bound, two-baryon bound states correspond to a superposition of $p - n$ and $\Delta - \Delta$ total spin $S = 1$ states, and also $\Delta - \Delta$, $S = 3$ states. The more weakly bound, bound states are associated with a superposition of $p - n$ and $\Delta - \Delta$ total spin $S = 0$ states and, in addition, $\Delta - \Delta$, $S = 2$ states. In contrast to the $I = 0$ states, we find that for the maximum isospin $I = 3$ sector there are also strongly bound and weakly bound bound states in the lowest spin sectors $S = 0, 1$ and *no* bound states if $S = 2, 3$. These results are in agreement with our previous results of Ref. [9] that the attraction between the two particles decreases with increasing I .

It is important to note that although our results are obtained using fairly complicated machinery, in the end, a simple picture emerges for the formation of baryon-baryon bound states. There is a correspondence between the lattice B-S equation, in relative coordinates, and a lattice resolvent operator Schrödinger equation. The two-baryon system in relative coordinates behaves approximately like that of a non-relativistic one-particle lattice Hamiltonian $T + V$ with lattice kinetic energy $-\kappa^3 \Delta_\ell$, where Δ_ℓ is the spatial lattice Laplacian and the potential energy V is $\kappa^2 V'$, the quasi-meson exchange space range-one potential which dominates the kinetic energy for small κ . The attractive or repulsive nature of the interaction depends on the isospin, spin spectral structure of V at a single site of space-range one. Because of the κ^3 dependence of the kinetic energy T and the κ^2 dependence of the potential energy, there is no minimal critical value of the interaction strength needed for the presence of a bound state. This is in contrast to the case of the Birman-Schwinger (see Ref. [28]) bound for a $-\Delta_\ell + \lambda V'$.

If V' is everywhere positive, there is a critical negative λ needed for a bound state to occur in three space dimensions.

A brief report on the results given in this paper appears in Ref. [29]. The present article is meant to be much more self-contained, describing in a relatively complete way the details about our methods and techniques. It is organized as follows. In Section 2, we mathematically define the model, describe the standard construction of the underlying physical Hilbert state $\mathcal{H}$, and introduce the fields which naturally emerge as those quantities associated with the physical baryon particles. To do this, we use a hyperplane decoupling expansion as done for quantum field theories in the continuum (see [30]), but adapted to the lattice. In Section 3, the one-particle analysis is done, and the baryon-baryon bound states are determined below the two-baryon threshold in Section 4. In order to keep the text as fluid as possible, trying to separate the main ideas of our methods from unavoidable sometimes heavy technical details, Sections 3 and 4 are subdivided according to their principal logical steps and details about specific points are given in the Appendices A-F. The similarity between the lattice B-S, in relative coordinates, and the lattice resolvent operator Schrödinger equation is exploited in Section 5. Finally, in Section 6, we end the paper with conclusions and prospective ideas, and a list of important open problems.

To close the Introduction, it is useful to make some comments on the seemingly awkward form of the two- and four- baryon correlation functions that we use in the sequel and which play an important role in our method. We explain the guiding principle for obtaining the form of correlation functions, which also holds in the continuum spacetime. Consider the physical Hilbert space matrix elements $(\chi, e^{-H|t|} \psi)$, where H is the energy operator, $e^{-H|t|}$ is the imaginary-time evolution operator (semigroup) and χ and ψ are both in the subspace of one- or two-baryon states. We have a F-K formula at our disposal and applying it to the two time orderings in t gives the form of the correlation functions. It is these correlation functions, that have appropriate imaginary-time decay properties, which allows our analysis to go through.

2. THE MODEL, THE PHYSICAL HILBERT SPACE AND BARYON FIELDS

We now introduce our $SU(3)$, $d + 1$ dimensional, imaginary-time lattice QCD model, where $d = 3$ is the space dimension. The partition function is given formally by $Z = \int e^{-S(\psi, \bar{\psi}, g)} d\psi d\bar{\psi} d\mu(g)$, and for a function $F(\bar{\psi}, \psi, g)$, the normalized correlations are denoted by

$$\langle F(\bar{\psi}, \psi, g) \rangle = \frac{1}{Z} \int F(\bar{\psi}, \psi, g) e^{-S(\psi, \bar{\psi}, g)} d\psi d\bar{\psi} d\mu(g). \quad (2.1)$$

The model action $S \equiv S(\psi, \bar{\psi}, g)$ is Wilson's action [1]

$$S = \frac{\kappa}{2} \sum_{\alpha, \beta} \bar{\psi}_{a, \alpha, f}(u) \Gamma_{\alpha\beta}^{\sigma e^\mu} (g_{u, u+\sigma e^\mu})_{ab} \psi_{b, \beta, f}(u + \sigma e^\mu) + \sum_{u \in \mathbb{Z}_o^4} \bar{\psi}_{a, \alpha, f}(u) M_{\alpha\beta} \psi_{a, \beta, f}(u) - \frac{1}{g_0^2} \sum_p \chi(g_p), \quad (2.2)$$

where, besides the sum over repeated indices $\alpha, \beta = 1, 2, 3, 4$ (spin), $a = 1, 2, 3$ (color) and $f = +1/2, -1/2 \equiv +, -$ (isospin), the first sum runs over $u = (u^0, \vec{u}) = (u^0, u^1, u^2, u^3) \in \mathbb{Z}_o^4 \equiv \{\pm 1/2, \pm 3/2, \pm 5/2, \dots\} \times \mathbb{Z}^3$, $\sigma = \pm 1$ and $\mu = 0, 1, 2, 3$. Here, we are adopting the label 0 for the time direction and e^μ , $\mu = 0, 1, 2, 3$, denotes the unit lattice vector for the μ -direction. The choice of the shifted lattice for the time direction, avoiding the zero-time coordinate, is so that, in the continuum limit, two-sided equal time limits of quark Fermi fields correlations can be accommodated. At each site $u \in \mathbb{Z}_o^4$, there are fermionic Grassmann fields $\psi_{a\alpha f}(u)$, associated with a quark, and fields $\bar{\psi}_{a\alpha f}(u)$, associated with an anti-quark, which carry a Dirac spin index α , an SU(3) color index a and isospin f . We refer to $\alpha = 1, 2$ as *upper* spin indices and $\alpha = 3, 4$ or $+$ or $-$ respectively, as *lower* ones. For each nearest neighbor oriented lattice bond $\langle u, u \pm e^\mu \rangle$ there is an SU(3) matrix $U(g_{u, u \pm e^\mu})$ parametrized by the gauge group element $g_{u, u \pm e^\mu}$ and satisfying $U(g_{u, u+e^\mu})^{-1} = U(g_{u+e^\mu, u})$. Associated with each lattice oriented plaquette p there is a plaquette variable $\chi(U(g_p))$ where $U(g_p)$ is the orientation-ordered product of matrices of SU(3) of the plaquette oriented bonds, and χ is the trace. For notational simplicity, we sometimes drop U from $U(g)$. Concerning the parameters, we take the quark-gauge coupling or hopping parameter $\kappa > 0$. Also, $g_0 > 0$ describes the pure gauge strength and $M \equiv M(m, \kappa) = (m + 2\kappa)I_4$, I_n being the $n \times n$ identity matrix. Given κ , for simplicity and without loss of generality, $m > 0$ is chosen such that $M_{\alpha\beta} = \delta_{\alpha\beta}$, meaning that $m + 2\kappa = 1$. Also, we take $\Gamma^{\pm e^\mu} = -I_4 \pm \gamma^\mu$, where γ_μ , satisfying $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}I_4$, are the 4×4 hermitian traceless anti-commuting Dirac matrices $\gamma^0 = \begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}$

and $\gamma^j = \begin{pmatrix} 0 & i\sigma^j \\ -i\sigma^j & 0 \end{pmatrix}$; and σ^j , $j = 1, 2, 3$, denotes the hermitian traceless anti-commuting 2×2 Pauli spin matrices. The measure $d\mu(g)$ is the product measure over non-oriented bonds of normalized SU(3) Haar measures (see [31]). There is only one integration variable per bond, so that g_{uv} and g_{vu}^{-1} are not treated as distinct integration variables. The integrals over Grassmann fields are defined according to [32]. For a polynomial in the Grassmann variables with coefficients depending on the gauge variables, the fermionic integral is defined as the coefficient of the monomial of maximum degree, i.e. of $\prod_{u, \ell} \bar{\psi}_\ell(u) \psi_\ell(u)$, $\ell \equiv (\alpha, a, f)$. In Eq. (2.1), $d\psi d\bar{\psi}$ means $\prod_{u, \ell} d\psi_\ell(u) d\bar{\psi}_\ell(u)$ such that, with a normalization $\mathcal{N}_1 = \langle 1 \rangle$, we have $\langle \psi_{\ell_1}(x) \bar{\psi}_{\ell_2}(y) \rangle =$

$(1/\mathcal{N}_1) \int \psi_{\ell_1}(x) \bar{\psi}_{\ell_2}(y) e^{-\sum_{u, \ell_3, \ell_4} \bar{\psi}_{\ell_3}(u) \mathcal{O}_{\ell_3 \ell_4} \psi_{\ell_4}(u)} d\psi d\bar{\psi} = \mathcal{O}_{\alpha_1, \alpha_2}^{-1} \delta_{a_1 a_2} \delta_{f_1 f_2} \delta(x - y)$, with a Kronecker delta for space-time coordinates, and where $\mathcal{O}$ is diagonal in the color and isospin indices.

Throughout the paper, we work in the strong coupling regime i.e. we take $0 < g_0^{-1} \ll \kappa \ll 1$, $m = 1 - 2\kappa \lesssim 1$. With this restriction on the parameters, and since we have chosen $\Gamma^{\pm e^\mu}$ matrices within the two-parameter family described in Ref. [13], positivity is preserved and there is a quantum mechanical Hilbert space $\mathcal{H}$ of physical states (see below). Furthermore, the condition $m > 0$ guarantees that the one-particle free Fermion dispersion curve increases in each positive momentum component and is convex for small momenta.

The action of Eq. (2.2) is invariant by the gauge transformations given by, for $x \in \mathbb{Z}_o^4$ and $h(x) \in \text{SU}(3)$,

$$\begin{aligned} \psi(x) &\mapsto h(x) \psi(x) \\ \bar{\psi}(x) &\mapsto \bar{\psi}(x) [h(x)]^{-1} \\ U(g_{x+e^\mu, x}) &\mapsto [h(x+e^\mu)]^{-1} U(g_{x+e^\mu, x}) h(x). \end{aligned} \quad (2.3)$$

Concerning the global SU(2) isospin symmetry of the action, we follow the treatment given in Ref. [10]. We treat isospin symmetry at the level of correlation functions (rather than as operators in the physical Hilbert space) in Appendix A. Other symmetries of the action in Eq. (2.2), such as time reversal, charge conjugation, parity, coordinate reflections and spatial rotations, which can be implemented by unitary (anti-unitary for time reversal) operators on the physical Hilbert space $\mathcal{H}$, were treated in Ref. [5] and do *not* affect the isospin indices. For $\kappa = 0$, we recover the spin SU(2) symmetry, separately in the lower and the upper spin indices. Furthermore, in Appendix B, a new time *reflection* symmetry, to be distinguished from time reversal, of the correlation functions is shown to be useful for obtaining additional relations between correlations.

By polymer expansion methods (see Refs. [13, 33]), the thermodynamic limit of correlations exists and truncated correlations have exponential tree decay. The limiting correlation functions are lattice translational invariant. Furthermore, the correlation functions extend to analytic functions in the coupling parameters κ and $\beta = 1/g_0^2$. For the formal hopping parameter expansion, see Refs. [2, 3].

The underlying quantum mechanical physical Hilbert space $\mathcal{H}$ and the EM operators H and P^j , $j = 1, 2, 3$ are defined as in Refs. [5, 6]. We start from gauge invariant correlations, with support restricted to $u^0 = 1/2$ and we let $T_0^{x^0}$, $T_i^{x^i}$, $i = 1, 2, 3$, denote translation of the functions of Grassmann and gauge variables by $x^0 \geq 0$, $\vec{x} = (x^1, x^2, x^3) \in \mathbb{Z}^3$. For F and G only depending on coordinates with $u^0 = 1/2$, we have the F-K formula

$$(G, T_0^{x^0} \check{T}_1^{x^1} \check{T}_2^{x^2} \check{T}_3^{x^3} F)_{\mathcal{H}} = \langle [T_0^{x^0} \vec{T} \vec{x} F] \Theta G \rangle, \quad (2.4)$$

where $T^{\vec{x}} = T_1^{x^1} T_2^{x^2} T_3^{x^3}$ and Θ is an anti-linear operator which involves time reflection. Following Ref. [13], the

action of Θ on single fields is given by

$$\begin{aligned}\Theta\bar{\psi}_{a\alpha}(u) &= (\gamma_0)_{\alpha\beta}\psi_{a\beta}(tu), \\ \Theta\psi_{a\alpha}(u) &= \bar{\psi}_{a\beta}(tu)(\gamma_0)_{\beta\alpha};\end{aligned}$$

where $t(u^0, \vec{u}) = (-u^0, \vec{u})$, for A and B monomials, $\Theta(AB) = \Theta(B)\Theta(A)$; and for a function of the gauge fields $\Theta f(\{g_{uv}\}) = f^*(\{g_{(tu)(tv)}\})$, $u, v \in \mathbb{Z}_o^{d+1}$, where $*$ means complex conjugate. Θ extends anti-linearly to the algebra. For simplicity, we do not distinguish between Grassmann, gauge variables and their associated Hilbert space vectors in our notation. As linear operators in $\mathcal{H}$, $\tilde{T}_\mu$, $\mu = 0, 1, 2, 3$, are mutually commuting; $\tilde{T}_0$ is self-adjoint, with $-1 \leq \tilde{T}_0 \leq 1$, and $\tilde{T}_{j=1,2,3}$ are unitary. So, we write $\tilde{T}_j = e^{iP^j}$ and $\vec{P} = (P^1, P^2, P^3)$ is the self-adjoint momentum operator. Its spectral points are $\vec{p} \in \mathbf{T}^3 \equiv (-\pi, \pi]^3$. Since $\tilde{T}_0^2 \geq 0$, the energy operator $H \geq 0$ can be defined by $\tilde{T}_0^2 = e^{-2H}$. We call a point in the EM spectrum associated with spatial momentum $\vec{p} = \vec{0}$ as a mass and, to be used below, we let $\mathcal{E}(\lambda^0, \vec{\lambda})$ be the product of the spectral families for the operators $\tilde{T}_0$, $\tilde{T}_1$, $\tilde{T}_2$ and $\tilde{T}_3$. By the spectral theorem (see Ref. [28], we have

$$\tilde{T}_0 = \int_{-1}^1 \lambda^0 dE_0(\lambda^0) \quad , \quad \tilde{T}_{j=1,2,3} = \int_{-\pi}^{\pi} \lambda^j dF_j(\lambda^j),$$

so that $\mathcal{E}(\lambda^0, \vec{\lambda}) = E_0(\lambda^0) \prod_{j=1}^3 F_j(\lambda^j)$. The positivity condition $\langle F\Theta F \rangle \geq 0$ is established in [13], but there may be nonzero F 's such that $\langle F\Theta F \rangle = 0$. If the collection of such F 's is denoted by $\mathcal{N}$, a pre-Hilbert space $\mathcal{H}'$ can be constructed from the inner product $\langle G\Theta F \rangle$ and the physical Hilbert space $\mathcal{H}$ is the completion of the quotient space $\mathcal{H}'/\mathcal{N}$, including also the cartesian product of the inner space sectors, the color space $\mathbb{C}^3$, the spin space $\mathbb{C}^4$ and the isospin space $\mathbb{C}^2$.

Here, we analyze the one and two-baryon sectors of $\mathcal{H}$. Points in the EM spectrum are detected as singularities in momentum space spectral representations of suitable two- and four-baryon field correlations given below.

3. ONE-PARTICLE SPECTRUM

In this section, we restrict our attention to the subspace $\mathcal{H}_o \subset \mathcal{H}$ generated by an odd number of $\tilde{\psi}$'s where the baryons lie (the tilde means the presence or absence of a bar). The mesons lie in the even subspace $\mathcal{H}_e$. We treat the one-baryon states adapting the methods of Refs. [5, 9].

3.1. Baryon fields and particle identifications

The decoupling of hyperplane method (see Refs. [5, 6]) reveals that the gauge invariant barred fields listed below create baryon particles. We point out that the same

method applies to QCD models with any number of flavors. In particular, for the SU(3) invariant three flavor case, the method would give the eightfold way. Here, the baryon particles are labeled by one-baryon total isospin (1st index), the one baryon z -component isospin (2nd index) and z -component of total spin (3rd index).

Let a hat $\hat{}$ mean the presence or absence of a bar simultaneously for *all* ψ 's. The decoupling of hyperplane expansion shows that the baryon fields are given by (with all fields evaluated at the same point)

$$\begin{cases} \hat{B}_{\frac{1}{2}\frac{1}{2}\frac{1}{2}} = \frac{\epsilon_{abc}}{3\sqrt{2}} (\hat{\psi}_{a\pm}\hat{\psi}_{b\mp} - \hat{\psi}_{a\pm}\hat{\psi}_{b\mp})\hat{\psi}_{c\pm}, \\ \hat{B}_{\frac{1}{2}\frac{-1}{2}\frac{1}{2}} = \frac{\epsilon_{abc}}{3\sqrt{2}} (\hat{\psi}_{a\pm}\hat{\psi}_{b\mp} - \hat{\psi}_{a\pm}\hat{\psi}_{b\mp})\hat{\psi}_{c\pm}, \end{cases} \quad (3.5)$$

$$\begin{cases} \hat{B}_{\frac{3}{2}\frac{1}{2}\frac{1}{2}} = \frac{\epsilon_{abc}}{6} (\hat{\psi}_{a++}\hat{\psi}_{b--} + 2\hat{\psi}_{a+-}\hat{\psi}_{b-+})\hat{\psi}_{c++}, \\ \hat{B}_{\frac{3}{2}\frac{1}{2}\frac{-1}{2}} = \frac{\epsilon_{abc}}{6} (2\hat{\psi}_{a++}\hat{\psi}_{b--} + \hat{\psi}_{a+-}\hat{\psi}_{b-+})\hat{\psi}_{c-+}, \\ \hat{B}_{\frac{3}{2}\frac{1}{2}\frac{3}{2}} = \frac{\epsilon_{abc}}{2\sqrt{3}} \hat{\psi}_{a\pm}\hat{\psi}_{b\pm}\hat{\psi}_{c\pm}, \end{cases} \quad (3.6)$$

$$\begin{cases} \hat{B}_{\frac{3}{2}\frac{-1}{2}\frac{1}{2}} = \frac{\epsilon_{abc}}{6} (2\hat{\psi}_{a++}\hat{\psi}_{b--} + \hat{\psi}_{a+-}\hat{\psi}_{b-+})\hat{\psi}_{c+-}, \\ \hat{B}_{\frac{3}{2}\frac{-1}{2}\frac{-1}{2}} = \frac{\epsilon_{abc}}{6} (\hat{\psi}_{a++}\hat{\psi}_{b--} + 2\hat{\psi}_{a+-}\hat{\psi}_{b-+})\hat{\psi}_{c--}, \\ \hat{B}_{\frac{3}{2}\frac{-1}{2}\frac{3}{2}} = \frac{\epsilon_{abc}}{2\sqrt{3}} \hat{\psi}_{a\pm}\hat{\psi}_{b\pm}\hat{\psi}_{c\pm}, \end{cases} \quad (3.7)$$

$$\begin{cases} \hat{B}_{\frac{3}{2}\frac{-3}{2}\frac{1}{2}} = \frac{\epsilon_{abc}}{2\sqrt{3}} \hat{\psi}_{a\pm}\hat{\psi}_{b\pm}\hat{\psi}_{c\mp}, \\ \hat{B}_{\frac{3}{2}\frac{-3}{2}\frac{3}{2}} = \frac{\epsilon_{abc}}{6} \hat{\psi}_{a\pm}\hat{\psi}_{b\pm}\hat{\psi}_{c\pm}, \end{cases} \quad (3.8)$$

$$\begin{cases} \hat{B}_{\frac{3}{2}\frac{3}{2}\frac{1}{2}} = \frac{\epsilon_{abc}}{2\sqrt{3}} \hat{\psi}_{a\pm}\hat{\psi}_{b\pm}\hat{\psi}_{c\mp}, \\ \hat{B}_{\frac{3}{2}\frac{3}{2}\frac{3}{2}} = \frac{\epsilon_{abc}}{6} \hat{\psi}_{a\pm}\hat{\psi}_{b\pm}\hat{\psi}_{c\pm}, \end{cases} \quad (3.9)$$

where the index α in $\hat{\psi}_{a,\alpha,f}$ is a *lower* spin component $\alpha = 3, 4 \equiv +, -$.

The baryon fields in Eqs. (3.5)-(3.9) obey the normalization $\langle B_i \bar{B}_j \rangle^{(0)} = -\delta_{ij}$, where i, j are collective indices and the superscript (0) denotes $\kappa = 0$ in the hopping term in the action S given in Eq. (2.2). Below, the superscript (n) means the *coefficient* of κ^n , $n = 0, 1, \dots$ We recall that, at $\kappa = 0$, our model also has a global SU(2) spin invariance in the space of lower and upper components.

A brief description of the hyperplane decoupling expansion method, and an important product structure for special third order κ derivatives, at $\kappa = 0$, for the two-baryon correlation function defined below is given in Appendix B [see Eq. (B.4)], together with the isospin identification for the above baryon fields with physical particles. The method reveals the fields and associated particles so that no guesswork or ansatz has to be used.

In analogy with the two-flavor SU(3) gauge QCD in the continuum with only the up (electric charge $+2/3$, z -component isospin $+1/2$) and the down (electric charge $-1/3$, z -component isospin $-1/2$) quarks, we identify the particles associated with the barred fields in Eqs. (3.5)-(3.9). The proton ($p_{\pm}$) and the neutron ($n_{\pm}$) are, respectively, described in Eq. (3.5). Δ^+ , Δ^0 and Δ^- are identified with the fields in Eqs. (3.6)-(3.8). Finally, Δ^{++} corresponds to the baryon fields in Eq. (3.9). The

corresponding z -component isospin conjugated particles are described by the unbarred fields.

We remark that, although not obvious, the fields in Eqs. (3.5)-(3.9) are the same as those obtained by the construction of gauge-invariant (colorless) wave functions as products of spin symmetric, flavor symmetric wave functions, except for the proton and neutron. For the proton and neutron, wave functions of Eq. (3.5) are the same as the sum of products of wave functions that are antisymmetric in both spin and flavor in the pairs 12, 13 and 23.

3.2. Two-baryon correlations and the one-particle analysis

The baryon-baryon correlation function is defined by (χ here denotes the characteristic function)

$$G_{\ell_1\ell_2}(u, v) = \langle B_{\ell_1}(u) \bar{B}_{\ell_2}(v) \rangle \chi_{u^0 \leq v^0} - \langle \bar{B}_{\ell_1}(u) B_{\ell_2}(v) \rangle^* \chi_{u^0 > v^0} \equiv G_{\ell_1\ell_2}(u - v), \quad (3.10)$$

where $\ell = (I, I_z, s)$ is a collective index. By isospin symmetry $G_{\ell_1\ell_2}(x = u - v)$ is diagonal in I, I_z and, for I fixed, independent of I_z . The consequences of isospin symmetry for the two-point functions, as well as for the four-point functions given below, are discussed in Appendix A, where important orthogonality relations and correlation function identities are derived. Also, the z -component of spin s is restricted to $\pm 1/2$, for $I = 1/2$ (for p and n) or $\pm 3/2, \pm 1/2$, for $I = 3/2$ (the Δ particles).

Setting $x^0 = u^0 - v^0 \neq 0$, we have the following spectral representation for G , with $\bar{B}_\ell \equiv \bar{B}_\ell(1/2, \vec{0})$,

$$G_{\ell_1\ell_2}(x) = - \int_{-1}^1 \int_{\mathbb{T}^3} (\lambda^0)^{|x^0|-1} e^{i\vec{\lambda} \cdot \vec{x}} \times d\lambda (\bar{B}_{\ell_1}, \mathcal{E}(\lambda^0, \vec{\lambda}) \bar{B}_{\ell_2})_{\mathcal{H}}; \quad (3.11)$$

for $x \in \mathbb{Z}^{d+1}$, $x^0 \neq 0$, and is an even function of $\vec{x}$.

Following Proposition 1 of Ref. [5], we verify how Eq. (3.11) is obtained. First, we derive a spectral representation for $u^0 < v^0$. With this time ordering, we have

$$G_{\ell_1\ell_2}(u, v) = \langle B_{\ell_1}(u) \bar{B}_{\ell_2}(v) \rangle = - \langle [T_0^{v^0 - u^0 - 1} \bar{T}^{\vec{v} - \vec{u}} \bar{B}_{\ell_2}(1/2, \vec{0})] B_{\ell_1}(-1/2, \vec{0}) \rangle.$$

But $B_\ell(-1/2, \vec{0}) = \Theta \bar{B}_\ell(1/2, \vec{0})$, so that

$$G_{\ell_1\ell_2}(u, v) = - \langle [T_0^{v^0 - u^0 - 1} \bar{T}^{\vec{v} - \vec{u}} \bar{B}_{\ell_2}(1/2, \vec{0})] \Theta \bar{B}_{\ell_1}(1/2, \vec{0}) \rangle.$$

By parity symmetry, $\bar{B}_\ell(w^0, \vec{0})$ has parity -1 so that $\vec{v} - \vec{u}$ can be replaced by $\vec{u} - \vec{v}$ here and below. The F-K formula of Eq. (2.4) gives the result. For $u^0 > v^0$, we write $G_{\ell_1\ell_2}(u, v) = - \langle [T_0^{u^0 - v^0 - 1} \bar{T}^{\vec{u} - \vec{v}} \bar{B}_{\ell_1}(-1/2, \vec{0})] \Theta \bar{B}_{\ell_2}(1/2, \vec{0}) \rangle^*$ and use the F-K formula. For $u^0 = v^0$ there is no spectral representation. By going backwards, the correlation of Eq. (3.10) is obtained from the Hilbert space inner product.

With this, upon separating the time zero contribution, the Fourier transform $\tilde{G}_{\ell_1\ell_2}(p) = \sum_{x \in \mathbb{Z}^4} G_{\ell_1\ell_2}(x) e^{-ip \cdot x}$ of $G_{\ell_1\ell_2}(x)$ admits the spectral representation

$$\tilde{G}_{\ell_1\ell_2}(p) = \tilde{G}_{\ell_1\ell_2}(\vec{p}) - (2\pi)^3 \int_{-1}^1 f(p^0, \lambda^0) d\lambda^0 \alpha_{\vec{p}, \ell_1\ell_2}(\lambda^0), \quad (3.12)$$

where

$$d\lambda^0 \alpha_{\vec{p}, \ell_1\ell_2}(\lambda^0) = \int_{\mathbb{T}^3} \delta(\vec{p} - \vec{\lambda}) d\lambda^0 d\vec{\lambda} (\bar{B}_{\ell_1}, \mathcal{E}(\lambda^0, \vec{\lambda}) \bar{B}_{\ell_2})_{\mathcal{H}}, \quad (3.13)$$

with $f(x, y) \equiv (e^{ix} - y)^{-1} + (e^{-ix} - y)^{-1}$, and we set $\tilde{G}(\vec{p}) = \sum_{\vec{x}} e^{-i\vec{p} \cdot \vec{x}} G(x^0 = 0, \vec{x})$.

The spectral representation given in Eq. (3.12) allows us to identify complex p singularities of $\tilde{G}(p)$ with points in the EM spectrum.

Using symmetries, $\tilde{G}_{\ell_1\ell_2}(\vec{p})$ is real diagonal and independent of isospin and color indices, diagonal and independent of the spin, at least up to order κ^6 .

From isospin symmetry, we see that the 20×20 two-point function matrix decomposes into blocks. There is one 2×2 block for the proton p and the same for the neutron n , and there are four 4×4 blocks associated to the Δ 's. Turning to the determination of the dispersion curves, as $G_{\ell_1\ell_2}$ is diagonal in I and I_z , and for fixed I , independent of I_z , we consider I and I_z fixed and denote the spin dependent two-point function by $G_{s_1 s_2}$. The dispersion curves $w(\vec{p}) \equiv w(\vec{p}, \kappa)$ are given by the zeroes of the determinant of $\tilde{\Gamma}_{s_1 s_2}(p)$, for p^0 in the positive imaginary axis, i.e.

$$\det \tilde{\Gamma}_{s_1 s_2}(p^0 = iw(\vec{p}), \vec{p}) = 0.$$

In Appendix B, we show that the 20×20 two-point function matrix $\tilde{\Gamma}_{s_1 s_2}(p)$ is *not* diagonal at $\vec{p} \neq \vec{0}$ (it is diagonal for $\vec{p} = \vec{0}$!). Thus, if we want to obtain the dispersion curves, this property prevents us to follow closely the analysis given in Ref. [5] for the mass spectrum. Using a different method, we show that there are 2 (4) dispersion curves, *not* necessarily distinct, for $I = 1/2$ ($3/2$), denoted by $w_j(\vec{p})$, $j = 1, 2$ for $I = 1/2$ and $j = 1, 2, 3, 4$ for $I = 3/2$; the masses m_j are given by $m_j = w_j(\vec{p} = \vec{0})$. The twenty approximate one-particle dispersion curves are identical up to and including $\mathcal{O}(\kappa^5)$ and, with

$$p_\ell^2 \equiv 2 \sum_{i=1}^3 (1 - \cos p^i), \quad (3.14)$$

are given by

$$w_j(\vec{p}) \equiv w_j(\vec{p}, \kappa) = [-3 \ln \kappa - 3\kappa^3/4 + p_\ell^2 \kappa^3/8] + r_j(\vec{p}), \quad (3.15)$$

where $r_j(\vec{p})$ is of $\mathcal{O}(\kappa^6)$, and its $r_j(\vec{p} = \vec{0})$ at $\mathcal{O}(\kappa^6)$ is zero. Thus, if there is a mass splitting, it is at least of $\mathcal{O}(\kappa^7)$. As also shown in Appendix B, for the case of 2×2 blocks associated with the proton and neutron, since the 2×2 matrices are diagonal by a parity-charge conjugation-time reflection symmetry, we can apply the

more powerful method of Ref. [5], which includes an application of the analytic implicit function theorem, to show that the corresponding (identical) $r_j(\vec{p})$ are analytic in κ and in each component $p^{i=1,2,3}$ of the momentum $\vec{p}$.

The spectral measure $d\lambda^0 \alpha_{\vec{p}, s_1 s_2}(\lambda^0)$ of Eq. (3.13) has the decomposition

$$d\lambda^0 \alpha_{\vec{p}, s_1 s_2}(\lambda^0) = \sum_j [Z_{j, s_1 s_2}(\vec{p}) \delta(\lambda^0 - e^{-w_j(\vec{p})})] d\lambda^0 + d\lambda^0 \nu_{s_1 s_2}(\lambda^0, \vec{p}),$$

where the first term corresponds to the one-particle contributions, and

$$Z_{j, s_1 s_2}(\vec{p})^{-1} = -(2\pi)^3 e^{w_j(\vec{p})} \frac{\partial \tilde{\Gamma}_{s_1 s_2}}{\partial \chi}(p^0 = i\chi, \vec{p})|_{\chi=w_j(\vec{p})}.$$

Let $\Gamma_{s_1 s_2}(u, v)$ denote the convolution inverse of the two-point function $G_{s_1 s_2}(u, v)$. We can show that $\Gamma_{s_1 s_2}(u, v)$ decays faster than $G_{s_1 s_2}(u, v)$, and hence the Fourier transform $\tilde{\Gamma}_{s_1 s_2}(p) = [\tilde{G}]_{s_1 s_2}^{-1}(p)$ of $\Gamma_{s_1 s_2}(x = u - v) \equiv \Gamma_{s_1 s_2}(u, v)$ has a larger region of analyticity in p^0 . Thus, the cofactor $[\text{cof } \tilde{\Gamma}]_{s_1 s_2}(p) / \det[\tilde{\Gamma}(p)] \equiv \tilde{\Gamma}_{s_1 s_2}^{-1}(p)$ provides a meromorphic extension of $\tilde{G}_{s_1 s_2}(p)$ up to near the two-particle threshold (precisely, the meson-baryon threshold, which is $\approx -5 \ln \kappa$), and the one-particle EM spectrum occurs, for each $\vec{p}$, as points given by the p^0 imaginary axis zeroes of $\det[\tilde{\Gamma}(p)]$. The two-point function convolution inverse $\Gamma(x)$ is defined by

$$\Gamma = (1 + G_d^{-1} G_n)^{-1} G_d^{-1}, \quad (3.16)$$

using a Neumann series, where G_d is the diagonal part of G given by

$$G_{d, s_1 s_2}(u, v) = G_{s_1 s_2}(u, u) \delta_{s_1 s_2} \delta_{uv}, \quad (3.17)$$

and G_n is the remainder

$$G_{n, s_1 s_2}(u, v) = G_{s_1 s_2}(u, v) - G_{d, s_1 s_2}(u, v). \quad (3.18)$$

By the decoupling of hyperplane method, we show the decay of G and Γ ; G_d^{-1} , G_n and Γ are bounded as matrix operators on $\ell_2(\mathbb{C}^{20} \times \mathbb{Z}_0^4)$; for small κ , G_n is small in norm so that the Neumann series in Eq. (3.16) converges. To obtain the results on the dispersion curves $w_j(\vec{p})$, up to order κ^6 , we need detailed information on the small distance behavior of $\Gamma_{s_1 s_2}$, which are obtained by showing explicit cancelations in the Neumann series and go beyond the hyperplane decoupling method (see Ref. [5] and Appendix B for more details).

We now give an intuitive argument for obtaining the mass and dispersion curves. The precise value requires a more refined analysis adapting the method used in Ref. [5], and depends on the short distance behavior of $\Gamma_{s_1 s_2}(x)$, which in turn depends on that of $G_{s_1 s_2}(x)$, via the Neumann series. By explicit calculation, $G_{s_1 s_2}(0) = -1$, $G_{s_1 s_2}(\pm e^0) = -\kappa^3$, $G_{s_1 s_2}(\pm e^j) = -\kappa^3/8$, $j = 1, 2, 3$, (correcting the misprinted values in Ref. [5]) such that

$$\tilde{G}_{s_1 s_2}(p) = [-1 - 2 \cos p^0 \kappa^3 - \frac{\kappa^3}{4} \sum_{j=1}^3 \cos p^j] \delta_{s_1 s_2} + \mathcal{O}(\kappa^6),$$

and then

$$\tilde{\Gamma}_{s_1 s_2}(p) = [-1 + 2 \cos p^0 \kappa^3 + \frac{\kappa^3}{4} \sum_{j=1}^3 \cos p^j] \delta_{s_1 s_2} + \mathcal{O}(\kappa^6).$$

Setting $\tilde{\Gamma}_{s_1 s_2}(p) = 0$ for $p^0 = i w_j(\vec{p})$, we obtain the dispersion curves of Eq. (3.15), which are shown to be the correct ones up to $\mathcal{O}(\kappa^5)$ by adapting the methods of Ref. [5].

Finally, with this, the λ^0 support of the measure $d\nu_{s_1 s_2}(\lambda^0, \vec{\lambda})$ is contained in $|\lambda^0| \leq |\kappa|^{5-\epsilon}$.

The decomposition of $\tilde{G}_{s_1 s_2}(p)$ is given by

$$\begin{aligned} \tilde{G}_{s_1 s_2}(p) &= \tilde{G}_{s_1 s_2}(\vec{p}) - \sum_j [Z_{j, s_1 s_2}(\vec{p}) \left[\frac{1}{e^{ip^0} - e^{w_j(\vec{p})}} \right. \\ &\quad \left. + \frac{1}{e^{-ip^0} - e^{w_j(\vec{p})}} \right] - (2\pi)^3 \int_{-1}^1 f(p^0, \lambda^0) \\ &\quad \times d\lambda^0 \nu_{\vec{p}, s_1 s_2}(\lambda^0)], \end{aligned} \quad (3.19)$$

4. TWO-BARYON SPECTRUM

In this section, we restrict our attention to the subspace $\mathcal{H}_e$ generated by an even number of $\tilde{\psi}$'s. We treat the two-baryon states adapting the methods of Ref. [9] (see also [29]).

4.1. Two-baryon states and four-baryon correlations in the individual and total spin basis

To determine baryon-baryon bound states, we consider the subspace of the subspace $\mathcal{H}_e \subset \mathcal{H}$ generated by the product of the twenty one-baryon fields given by $\bar{B}_{\ell_1}(\vec{x}_1) \bar{B}_{\ell_2}(\vec{x}_2)$. If we ignore possible linear dependencies, this space has dimension 400, which -upon using isospin symmetry- decomposes into two-baryon total isospin $I = 0, 1, 2, 3$ sectors, of dimensions 20, 108, 160 and 112, respectively. In this subsection, to reduce the complexity of the analysis, here we first restrict our attention to the $I = 0$ sector, where we expect to find a deuteron, $p - n$ bound state. The $I = 3$ states will be treated in Section 4.4. The more algebraically complex $I = 1, 2$ sectors are presently being analyzed [34]. The isospin orthogonality relations and correlation identities used here are derived in Appendix A.

The $I = 0$ states are given by the isospin Clebsch-Gordan (C-G) linear combinations

$$\bar{\tau}_{s_1 s_2}^{1/2}(x_1, x_2) = \sum_{I_1 + I_2 = 0} c_{I_1 I_2}^{\frac{1}{2}} \bar{B}_{\frac{1}{2}, I_1, s_1}(x_1) \bar{B}_{\frac{1}{2}, I_2, s_2}(x_2),$$

for the coupling of two isospin 1/2 baryons, and similarly, with $(1/2 \rightarrow 3/2, s_i \rightarrow t_i)$, for the coupling of two 3/2 isospins. We use $s_j (t_j)$ for isospin 1/2 (3/2) baryons and recall that the 1/2, 3/2 coupling does *not* give $I = 0$; and

we refer to this description of the two-particle states as the individual spin basis.

It turns out that the dominant interaction between baryons admits a simpler description in terms of total spin. The total spin states are obtained by taking C-G combinations of the τ 's. Namely,

$$\bar{T}_{SSz}^{1/2}(x_1, x_2) = \sum_{s_1+s_2=S_z} a_{s_1 s_2}^{1/2} \bar{\tau}_{s_1, s_2}^{1/2}(x_1, x_2), \quad (4.20)$$

and similarly, with $(1/2 \rightarrow 3/2, S \rightarrow T, s_i \rightarrow t_i)$, where the a 's are the spin C-G coefficients. Using their properties, we find that

$$\begin{aligned} \bar{T}_{SSz}^{1/2}(x_1, x_2) &= (-1)^{S+1} \bar{T}_{SSz}^{1/2}(x_2, x_1), \\ \bar{T}_{TZ}^{3/2}(x_1, x_2) &= (-1)^{T+1} \bar{T}_{TZ}^{3/2}(x_2, x_1), \end{aligned} \quad (4.21)$$

so that, for coincident points, the states vanish for $S = 0$, and for $T = 0, 2$, a kinematic statistical effect. The states are in the space symmetric (anti-symmetric) for $S = 1$, $T = 1, 3$ ($S = 0, T = 0, 2$).

We order the individual spin pairs as $(s_1, s_2) = (\frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, -\frac{1}{2}), (-\frac{1}{2}, \frac{1}{2}), (-\frac{1}{2}, -\frac{1}{2}); (t_1, t_2) = (\frac{3}{2}, \frac{3}{2}), (\frac{3}{2}, \frac{1}{2}), (\frac{3}{2}, -\frac{1}{2}), (\frac{3}{2}, -\frac{3}{2}), (\frac{1}{2}, \frac{3}{2}), \dots, (\frac{1}{2}, -\frac{3}{2}), (-\frac{1}{2}, \frac{3}{2}), \dots, (-\frac{1}{2}, -\frac{3}{2}), (-\frac{3}{2}, \frac{3}{2}), \dots, (-\frac{3}{2}, -\frac{3}{2})$, and the total spin pairs as $(S, S_z) = (1, 1), (T, T_z) = (1, 1), (S, S_z) = (1, 0), (T, T_z) = (1, 0), (S, S_z) = (1, -1), (T, T_z) = (1, -1), (T, T_z) = (3, 3), (3, 2), \dots, (3, -3), (S, S_z) = (0, 0), (T, T_z) = (0, 0), (T, T_z) = (2, 2), (2, 1), \dots, (2, -2)$. From Eq. (4.21) and in the total spin basis we see that the two baryon states are space symmetric (anti-symmetric) in the 1st through the 13th (last 7) components.

The individual and the total spin basis, in the $I = 0$ sector, are related by a real orthogonal transformation whose entries are C-G coefficients. Let $\mathcal{A} = [\mathcal{A}]_{i,j=1,\dots,20}$ be the 20×20 transformation matrix from the individual to the total spin basis. The nonzero elements of $\mathcal{A}$ are: $\mathcal{A}_{11} = 1 = \mathcal{A}_{45} = \mathcal{A}_{75} = \mathcal{A}_{13\ 20}$; $\mathcal{A}_{27} = \sqrt{3/10} = \mathcal{A}_{2\ 13} = \mathcal{A}_{6\ 12} = \mathcal{A}_{6\ 18}$; $\mathcal{A}_{2\ 10} = -\sqrt{2/5} = \mathcal{A}_{6\ 15}$; $\mathcal{A}_{32} = \sqrt{1/2} = \mathcal{A}_{33} = \mathcal{A}_{86} = \mathcal{A}_{89} = \mathcal{A}_{12\ 16} = \mathcal{A}_{12\ 19} = \mathcal{A}_{14\ 2} = \mathcal{A}_{16\ 6} = \mathcal{A}_{17\ 7} = \mathcal{A}_{19\ 12} = \mathcal{A}_{20\ 16}$; $\mathcal{A}_{48} = \sqrt{9/20} = \mathcal{A}_{4\ 17} = \mathcal{A}_{10\ 11} = \mathcal{A}_{10\ 14}$; $\mathcal{A}_{4\ 11} = -\sqrt{1/20} = \mathcal{A}_{4\ 14}$; $\mathcal{A}_{97} = \sqrt{1/5} = \mathcal{A}_{9\ 13} = \mathcal{A}_{11\ 12} = \mathcal{A}_{11\ 18}$; $\mathcal{A}_{9\ 10} = \sqrt{3/5} = \mathcal{A}_{11\ 15}$; $\mathcal{A}_{10\ 8} = \sqrt{1/20} = \mathcal{A}_{10\ 17}$; $\mathcal{A}_{14\ 3} = -\sqrt{1/2} = \mathcal{A}_{16\ 9} = \mathcal{A}_{17\ 13} = \mathcal{A}_{19\ 18} = \mathcal{A}_{20\ 19}$; $\mathcal{A}_{15\ 8} = 1/2 = \mathcal{A}_{15\ 14} = \mathcal{A}_{18\ 8} = \mathcal{A}_{18\ 11}$ and, finally, $\mathcal{A}_{15\ 11} = -1/2 = \mathcal{A}_{15\ 17} = \mathcal{A}_{18\ 14} = \mathcal{A}_{18\ 17}$.

4.2. Four-baryon correlations and two-baryon spectrum

As we do not know a priori which linear combination of states corresponds to a bound state, we consider the 20×20 matrix of (truncated) four-baryon functions

$$M(x_1, x_2; x_3, x_4) = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix} (x_1, x_2; x_3, x_4),$$

where, in the individual spin basis, and for $x_1^0 = x_2^0, x_3^0 = x_4^0, x_1^0 \leq x_3^0$, we have $M_{11; s_1 s_2 s_3 s_4}(x_1, x_2; x_3, x_4) = \langle \tau_{s_1 s_2}^{1/2}(x_1, x_2) \bar{\tau}_{s_3 s_4}^{1/2}(x_3, x_4) \rangle$, $M_{12; s_1 s_2 t_3 t_4}(x_1, x_2; x_3, x_4) = \langle \tau_{s_1 s_2}^{1/2}(x_1, x_2) \bar{\tau}_{t_3 t_4}^{3/2}(x_3, x_4) \rangle$, $M_{21; t_1 t_2 s_3 s_4}(x_1, x_2; x_3, x_4) = \langle \tau_{t_1 t_2}^{3/2}(x_1, x_2) \bar{\tau}_{s_3 s_4}^{1/2}(x_3, x_4) \rangle$ and $M_{22; t_1 t_2 t_3 t_4}(x_1, x_2; x_3, x_4) = \langle \tau_{t_1 t_2}^{3/2}(x_1, x_2) \times \bar{\tau}_{t_3 t_4}^{3/2}(x_3, x_4) \rangle$; and similarly for the total spin basis. For $x_1^0 > x_3^0$, M_{ij} is defined by interchanging the barred and unbarred fields and taking the complex conjugate. As in Ref. [9], $x_1, \dots, x_4$ are now taken in $\mathbb{Z}^4$.

M_{ij} has spectral representations which we use to detect bound states below the two-baryon threshold. We now obtain a spectral representation for $M_{ij; s_1 s_2 s_3 s_4}$. To simplify and make the analysis more transparent, we derive a spectral representation in terms of the basic baryon fields B_ℓ . By simply writing the suitable linear combinations of these representations, we obtain the spectral representations for the four-point functions of the τ_ℓ fields.

The starting point employs the F-K formula which furnishes a 4-point correlation function in terms of B_ℓ 's. For $x^0 \neq 0$, with $x = (x^0 = v^0 - u^0, \vec{x}) \in \mathbb{Z}^4$,

$$\begin{aligned} &(\bar{B}_{\ell_1}(1/2, \vec{u}_1) \bar{B}_{\ell_2}(1/2, \vec{u}_2), (\check{T}_0)^{|x^0|-1} \\ &\times \bar{T}^{\vec{x}} \bar{B}_{\ell_3}(1/2, \vec{u}_3) \bar{B}_{\ell_4}(1/2, \vec{u}_4))_{\mathcal{H}} = -\mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(x), \end{aligned}$$

where $\mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(x) \equiv \mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(u_1, u_2, u_3 + \vec{x}, u_4 + \vec{x})$ and, for $u_1^0 = u_2^0 = u^0$ and $u_3^0 = u_4^0 = v^0$,

$$\begin{aligned} \mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(u_1, u_2, u_3, u_4) &= \\ &\langle B_{\ell_1}(u_1) B_{\ell_2}(u_2) \bar{B}_{\ell_3}(u_3) \bar{B}_{\ell_4}(u_4) \rangle \chi_{u^0 \leq v^0} \\ &+ \langle \bar{B}_{\ell_1}(u_1) \bar{B}_{\ell_2}(u_2) B_{\ell_3}(u_3) B_{\ell_4}(u_4) \rangle^* \chi_{u^0 > v^0}. \end{aligned}$$

We obtain a spectral representation for $\mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(x)$ and its Fourier transform $\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k)$ which allows us to relate the complex k singularities of $\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k)$ to the EM spectrum. Inserting the spectral representations for $\check{T}_0$ and $\bar{T}_{i=1,2,3}$, and taking the Fourier transform gives

$$\begin{aligned} \tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k) &= \tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{k}) - (2\pi)^3 \int_{-1}^1 \int_{\mathbf{T}^3} f(k^0, \lambda^0) \\ &\times \delta(\vec{k} - \vec{\lambda}) d\lambda^0 d\vec{\lambda} (\bar{B}_{\ell_1}(1/2, \vec{u}_1) \bar{B}_{\ell_2}(1/2, \vec{u}_2), \\ &\times \mathcal{E}(\lambda^0, \vec{\lambda}) \bar{B}_{\ell_3}(1/2, \vec{u}_3) \bar{B}_{\ell_4}(1/2, \vec{u}_4))_{\mathcal{H}}, \end{aligned}$$

where $\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{k}) = \sum_{\vec{x} \in \mathbf{T}^3} e^{-i\vec{k} \cdot \vec{x}} \mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(x^0 = 0, \vec{x})$. The singularities of $\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k)$, for $k = (k^0 = i\chi, \vec{k} = 0)$ and $e^{\pm\chi} \leq 1$, are points in the mass spectrum.

To analyze $\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k)$, we follow the method of analysis for spin models as in Ref. [26]. To use a notation closer to that of Ref. [26], we relabel the time direction coordinates in $\mathcal{G}_{\ell_1 \ell_2 \ell_3 \ell_4}(x)$ by integer labels, with $u_i^0 - 1/2 = x_i^0, \vec{u}_i = \vec{x}_i, i = 1, \dots, 4$, and denote by $D_{\ell_1 \ell_2 \ell_3 \ell_4}$ the shifted correlation functions. Next, we write $D_{\ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3 + \vec{x}, x_4 + \vec{x})$, $x_1^0 = x_2^0$ and $x_3^0 = x_4^0$, $x^0 = x_3^0 - x_2^0$, where x_i and x are points on the $\mathbb{Z}^4$ lattice. Now, by translation invariance, we pass to difference coordinates and then to lattice relative coordinates

$\xi = x_2 - x_1$, $\eta = x_4 - x_3$ and $\tau = x_3 - x_2$ (see Figure 1 below). These lattice relative coordinates are a substitute for the usual center of mass and relative coordinates customarily used in the continuum. After doing this, we obtain

$$\begin{aligned} D_{\ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3 + \vec{x}, x_4 + \vec{x}) \\ = D_{\ell_1 \ell_2 \ell_3 \ell_4}(0, x_2 - x_1, x_3 - x_1 + \vec{x}, x_4 - x_1 + \vec{x}) \\ \equiv D_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{\xi}, \vec{\eta}, \tau + \vec{x}), \end{aligned}$$

and

$$\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k) = e^{i\vec{k} \cdot \vec{\tau}} \hat{D}_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{\xi}, \vec{\eta}, k),$$

where $\hat{D}_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{\xi}, \vec{\eta}, k) = \sum_{\tau \in \mathbb{Z}^4} D_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{\xi}, \vec{\eta}, \tau) e^{-ik \cdot \tau}$. Explicitly, we have

$$\begin{aligned} D_{\ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3, x_4) = & \langle B_{\ell_1}(x_1^0 + 1/2, \vec{x}_1) \\ & \times B_{\ell_2}(x_2^0 + 1/2, \vec{x}_2) \bar{B}_{\ell_3}(x_3^0 + 1/2, \vec{x}_3) \\ & \times \bar{B}_{\ell_4}(x_4^0 + 1/2, \vec{x}_4) \rangle \chi_{x_2^0 \leq x_3^0} \\ & + \langle \bar{B}_{\ell_1}(x_1^0 + 1/2, \vec{x}_1) \bar{B}_{\ell_2}(x_2^0 + 1/2, \vec{x}_2) \\ & \times B_{\ell_3}(x_3^0 + 1/2, \vec{x}_3) B_{\ell_4}(x_4^0 + 1/2, \vec{x}_4) \rangle^* \chi_{x_2^0 > x_3^0}. \end{aligned} \quad (4.22)$$

The point of the considerations above is that the singularities of $\tilde{\mathcal{G}}_{\ell_1 \ell_2 \ell_3 \ell_4}(k)$ are the same as those of $\hat{D}_{\ell_1 \ell_2 \ell_3 \ell_4}(\vec{\xi}, \vec{\eta}, k)$.




Figure 1: Lattice relative coordinates.

Analogously, reinstating the linear combinations, for $M_{\ell_1 \ell_2 \ell_3 \ell_4}$, we obtain the spectral representation

$$\begin{aligned} \tilde{M}_{\ell_1 s_2 s_3 s_4}(k) = & \tilde{M}_{s_1 s_2 s_3 s_4}(\vec{k}) - (2\pi)^3 \int_{-1}^1 \int_{\mathbf{T}^3} f(k^0, \lambda^0) \\ & \times \delta(\vec{k} - \vec{\lambda}) d\lambda^0 d\vec{\lambda} (\bar{\tau}_{s_1}(1/2, \vec{u}_1) \bar{\tau}_{s_2}(1/2, \vec{u}_2), \\ & \times \mathcal{E}(\lambda^0, \vec{\lambda}) \bar{\tau}_{s_3}(1/2, \vec{u}_3) \bar{\tau}_{s_4}(1/2, \vec{u}_4))_{\mathcal{H}}, \end{aligned}$$

where, for $x^0 \neq 0$, and with $x = (x^0 = v^0 - u^0, \vec{x}) \in \mathbb{Z}^4$,

$$\tilde{M}_{s_1 s_2 s_3 s_4}(\vec{k}) = \sum_{\vec{x} \in \mathbb{Z}^3} e^{-i\vec{k} \cdot \vec{x}} M_{s_1 s_2 s_3 s_4}(x^0 = 0, \vec{x}).$$

The singularities of $\tilde{M}_{s_1 s_2 s_3 s_4}$, for $k = (k^0 = i\chi, \vec{k} = \vec{0})$ and $e^{\pm\chi} \leq 1$ are points in the mass spectrum. To determine $\tilde{M}_{s_1 s_2 s_3 s_4}$ we employ a lattice version of the B-S equation, as discussed below. To detect baryon-baryon bound states below the lowest two-baryon threshold, we write $k^0 = i(2\bar{m} - \epsilon)$, where $\bar{m}$ is the smallest baryon mass and determine the value of $\epsilon > 0$ that gives the singularity. We call this special value of ϵ the binding energy of the corresponding bound state.

4.3. The B-S equation, the ladder approximation and $I = 0$ bound states

Next, we write a B-S equation for M as,

$$M = M_0 + M_0 K M, \quad (4.23)$$

and in contrast to the continuum case, we use what we call an equal time representation.

In terms of kernels, with $x_1^0 = x_2^0$ and $x_3^0 = x_4^0$, we have

$$\begin{aligned} M(x_1, x_2, x_3, x_4) = & M_0(x_1, x_2, x_3, x_4) + \int M_0(x_1, x_2, y_1, y_2) \\ & \times K(y_1, y_2, y_3, y_4) M(y_3, y_4, x_3, x_4) \\ & \times \delta(y_1^0 - y_2^0) \delta(y_3^0 - y_4^0) dy_1 dy_2 dy_3 dy_4, \end{aligned}$$

where we use a continuum notation for sums over lattice points. $K(y_1, y_2, y_3, y_4)$ is called the B-S kernel. We remark that in the imaginary time continuum quantum field theory formulation with Euclidean invariance all the B-S equation coordinates are taken to run over all space-time.

Formally, this B-S equation defines $K = M_0^{-1} - M^{-1}$. Our strategy is to compute K in the leading κ order, which we call a ladder approximation L to K , and then solve the B-S equation for M using L . In the B-S equation, M_0 is obtained from M by erroneously applying Wick's theorem to the one-baryon composite fields $\bar{B}$ in M such as,

$$\begin{aligned} -\langle B_{\ell_1}(x_1) \bar{B}_{\ell_3}(x_3) \rangle \langle B_{\ell_2}(x_2) \bar{B}_{\ell_4}(x_4) \rangle \\ + \langle B_{\ell_1}(x_1) \bar{B}_{\ell_4}(x_4) \rangle \langle B_{\ell_2}(x_2) \bar{B}_{\ell_3}(x_3) \rangle, \end{aligned}$$

for the term $\langle B_{\ell_1}(x_1) B_{\ell_2}(x_2) \bar{B}_{\ell_3}(x_3) \bar{B}_{\ell_4}(x_4) \rangle$ in $M_{\ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3, x_4)$. With this, we obtain

$$\begin{aligned} F_{0, \ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3, x_4) \equiv & -G_{\ell_1 \ell_3}(x_1, x_3) G_{\ell_2 \ell_4}(x_2, x_4) \\ & + G_{\ell_1 \ell_4}(x_1, x_4) G_{\ell_2 \ell_3}(x_2, x_3). \end{aligned} \quad (4.24)$$

By referring to the definition of the two-baryon function $G_{\ell_1 \ell_4}$ of Eq. (3.10), we see that Eq. (4.24) holds for all temporal orderings.

In terms of the $(\vec{\xi}, \vec{\eta}, \tau)$ lattice relative coordinates and taking the Fourier transform in τ only, the B-S equation becomes (see [26])

$$\begin{aligned} \hat{M}(\vec{\xi}, \vec{\eta}, k) = & \hat{M}_0(\vec{\xi}, \vec{\eta}, k) + \int \hat{M}_0(\vec{\xi}, \vec{\xi}', k) \\ & \times \hat{K}(-\vec{\xi}', -\vec{\eta}', k) \hat{M}(\vec{\eta}', \vec{\eta}, k) d\vec{\xi}' d\vec{\eta}'. \end{aligned}$$

The above B-S equation corresponds, roughly speaking, to an equation for the kernel of a one-particle lattice Schrödinger operator resolvent equation. The B-S kernel $\hat{K}(-\vec{\xi}', -\vec{\eta}', k)$, in general, acts as an energy-dependent, non-local potential (see Refs. [9, 10]). $\hat{M}_0$ is like the resolvent of the dressed noninteracting particles and $\hat{M}$ is the interacting resolvent. To facilitate the analysis, it is desirable to have $\hat{K}$ with the shortest range as possible. For example, in weakly coupled scalar models, with local interactions and in the weak coupling regime, as well

as in other spin and stochastic systems, the dominant contribution to $\hat{K}$ is local, and the ladder approximation dominates in the full model (see Refs. [26, 27, 35]). In general, the hyperplane decoupling method is used to obtain decay properties of K , which allows us to control the solution to the B-S equation.

To make sense of the B-S equation and the kernel K , we now define the space where M and M_0 act as matrix operators, which is given by $\ell'_2 = \ell'_2 \oplus \ell'_2$, such that $f = f_1 + f_2 \in \ell'_2$, f_1 (f_2) is associated with the first 13 (last 7) components; $f_1 \in \ell'_2(A_s)$, $f_2 \in \ell'_2(A_a)$, where s (a) means the symmetric (antisymmetric) subspace and

$$A_s = \{(x_1, x_2; j) \in \mathbb{Z}^4 \times \mathbb{Z}^4 \times (1, 2, \dots, 13) | x_1^0 = x_2^0\},$$

and

$$A_a = \{(x_1, x_2; j) \in \mathbb{Z}^4 \times \mathbb{Z}^4 \times (14, \dots, 20) | x_1^0 = x_2^0, \vec{x}_1 \neq \vec{x}_2\}.$$

Using the orthogonality properties of the two-point functions, in the total spin basis, we have

$$M_{0;S,S_z,S',S'_z}^{(0)}(x_1, x_2; x_3, x_4) = \delta_{SS'} \delta_{S_z S'_z} [-\delta(x_1 - x_3) \times \delta(x_2 - x_4) + (-1)^S \times \delta(x_1 - x_4) \delta(x_2 - x_3)],$$

and the same for $M_{0;T,T_z,T',T'_z}^{(0)}$, $T = 0, 1, 2, 3$. Hence, $M_0^{(0)}$ acts as $-2\mathbb{1} \neq 0$ on ℓ'_2 , and M_0 is invertible. To define the B-S operator $K = M_0^{-1} - M^{-1}$, we also need M^{-1} . For $M(x_1, x_2, x_3, x_4)$, setting

$$M(0) \equiv M(x, x, x, x) = M(0, 0, 0, 0),$$

we find that the κ^0 order is given by

$$M^{(0)}(x_1, x_2, x_3, x_4) = M^{(0)}(0) \delta(x_1 - x_2) \delta(x_1 - x_3) \times \delta(x_1 - x_4) + M_0^{(0)} [1 - \delta(x_1 - x_2) \delta(x_1 - x_3) \delta(x_1 - x_4)],$$

such that $M^{(0)}$ agrees with $M_0^{(0)}$, except at coincident points. Hence, to show M is invertible we need the invertibility of the matrix $M^{(0)}(0)$ which we analyze next.

By a lengthy computation given in Appendix C, we find that, in the total spin basis,

$$M^{(0)}(0) = M_1^{(0)}(0) \oplus M_1^{(0)}(0) \oplus M_1^{(0)}(0) \oplus M_b^{(0)}(0), \quad (4.25)$$

where

$$M_1^{(0)}(0) = -\frac{1}{9} \begin{pmatrix} 20 & -8\sqrt{5} \\ -8\sqrt{5} & 16 \end{pmatrix}, \quad (4.26)$$

and $M_b^{(0)}(0)$ acts as $-4\mathbb{1}$ on the components with indices from 7 to 13. The eigenvalues of $M_1^{(0)}(0)$ are -4 and 0 . The null space prohibits the invertibility of $M^{(0)}(0)$.

The zero eigenvector is $\frac{1}{3} \begin{pmatrix} 2 \\ \sqrt{5} \end{pmatrix}$ and the first line of the eigenvalue equation is

$$\langle \mathcal{T}_{S=1, S_z} [\mathcal{T}_{S=1, S_z} + \frac{\sqrt{5}}{2} \mathcal{T}_{T=1, T_z}] \rangle^{(0)} = 0.$$

The vanishing of this average suggests that there is a pointwise linear dependence relation at coincident points. Indeed, this turns out to be the case, and is proven in Appendix C that, at coincident points,

$$\mathcal{T}_{S=1, S_z} + \frac{\sqrt{5}}{2} \mathcal{T}_{T=1, T_z} = 0, \quad (4.27)$$

for $S_z = T_z = -1, 0, 1$.

The truncated four-point correlation M still has a null space in the symmetric subspace of ℓ'_2 . For this reason, we are forced to reduce the symmetric subspace for coincident points only, and we take the restriction of $M^{(0)}(0)$ to the orthogonal complement of the null space in the symmetric subspace. So, for coincident points we have a 10 dimensional space. For non-coincident points, $M^{(0)}$ agrees with $M_0^{(0)} \neq 0$, so that K is well-defined.

With this definition of K , and using its decay properties shown in Appendix D, there is a local potential, and energy-dependent space range-zero potential, as well as an energy-independent range-one potential. As in Ref. [9], we modify M at coincident points by a multiplicative constant $1/2$. Precisely, we replace $M(x_1, x_2, x_3, x_4)$ by

$$M'(x_1, x_2, x_3, x_4) = h(x_1, x_2) M(x_1, x_2, x_3, x_4) h(x_3, x_4),$$

where

$$h(x, y) = [1 - \delta(x - y)] + \frac{1}{\sqrt{2}} \delta(x - y),$$

and the B-S equation becomes

$$M' = M_0 + M_0 K' M'. \quad (4.28)$$

The τ variable Fourier transforms $\hat{M}$ and $\hat{M}'$ clearly have the same singularities. For simplicity of notation, we drop the prime and keep using M , $\hat{M}$, K and $\hat{K}$.

Next, to define our ladder approximation, we write a Neumann expansion representation for $K = M_0^{-1} - M^{-1}$. With

$$\delta M = M - M^{(0)} \quad , \quad \delta M_0 = M_0 - M_0^{(0)},$$

we have

$$K = \sum_{n=0}^{\infty} (-1)^n [(M_0^{(0)-1} \delta M_0)^n M_0^{(0)-1} - (M^{(0)-1} \delta M)^n M^{(0)-1}]. \quad (4.29)$$

For small κ , the dominant contributions to K come from coincident points at $\kappa = 0$ and a space range-one potential of order κ^2 . As discussed in Appendix E, it turns out that, because of cancelations $K^{(0)}(0) = (M_0^{(0)})^{-1}(0) - (M^{(0)})^{-1}(0) = 0$. Surprisingly, there is an attractive gauge field correlation effect which is responsible for this result. More explicitly, as a contribution to the definition of $h(x, y)$, we must consider the effect of a gauge integral over six overlapping bonds with the same orientation, which is given by $\mathcal{I}_6$ in Eq. (D6). Moreover, it is worth mentioning that if these six gauge fields are

uncorrelated three by three, meaning that $\mathcal{I}_6$ is replaced by $(\mathcal{I}_3)^2$, $\mathcal{I}_3$ being the integral over three overlapping bonds with the same orientation [see Eq. (B.1)], the associated attractive contribution to $K^{(0)}(0)$ becomes repulsive, leading to an overall repulsive $K^{(0)}(0)$.

Going further in the κ expansion, since δM is $\mathcal{O}(\kappa^2)$ and δM_0 is $\mathcal{O}(\kappa^3)$, we find that the dominant contribution to K is given by the $n = 1$ term in the Neumann expansion of Eq. (4.29). This defines our ladder approximation to K and is given by

$$L \equiv K^{(2)}\kappa^2 = (M^{(0)})^{-1}(\delta M)^{(2)}(M^{(0)})^{-1}\kappa^2. \quad (4.30)$$

To obtain an explicit formula for the kernel of L of Eq. (4.30), first we consider the contribution to L coming from the non-oriented geometric bond linking the sites 0 and e^1 . It is given by

$$\begin{aligned} K^{(2)}(0, e^1; 0, e^1) + K^{(2)}(0, e^1; e^1, 0) &= (M^{(0)})^{-1}(0, e^1; 0, e^1) \\ &\times M^{(2)}(0, e^1; 0, e^1)(M^{(0)})^{-1}(0, e^1; 0, e^1) \\ &+ (M^{(0)})^{-1}(0, e^1; e^1, 0)M^{(2)}(0, e^1; e^1, 0) \\ &\times (M^{(0)})^{-1}(0, e^1; e^1, 0). \end{aligned}$$

Since $M^{(0)} = M_0^{(0)} = -2 \mathbb{1}$, for non-coincident points, each $(M^{(0)})^{-1}$ factor can be replaced by $-1/2$. In the total spin basis, taking into account the space coordinate exchange symmetry (anti-symmetry) of $\mathcal{T}$ for $S = 1, T = 1, 3$ ($S = 0, T = 0, 2$) given in Eq. (4.21), the second term is the same as the first one.

The contribution $M^{(2)}(0, e^1; 0, e^1)\kappa^2$ is obtained by expanding the hopping term exponential in the action appearing in the numerator, which brings down a pair of oppositely oriented overlapping gauge field bounds. The gauge field integral of Eq. (D7) is performed using the Peter-Weyl theorem (see Ref. [2] for other more complicated and useful gauge integrals and Appendix D), and gives a factor $1/3$. In this way, the 11 element $M^{(2)}(0, e^1; 0, e^1)$ of the matrix $M^{(2)}$ is

$$M_{11, s_1 s_2 s_3 s_4}^{(2)}(0, e^1; 0, e^1) = \frac{1}{12} v_{s_1 s_2 s_3 s_4}^{(2)},$$

where, in the individual spin basis,

$$\begin{aligned} v_{s_1, s_2, s_3, s_4}^{(2)} &= \sum' c_{I_1 I_2}^{\frac{1}{2} \frac{1}{2}} c_{I_3 I_4}^{\frac{1}{2} \frac{1}{2}} \times \\ &\langle B_{\frac{1}{2} I_1 s_1} \bar{B}_{\frac{1}{2} I_3 s_3} \bar{\psi}_{a_1 \alpha_1 f_1} \psi_{a_1 \beta_2 f_2} \rangle^{(0)} \times \\ &\langle B_{\frac{1}{2} I_2 s_2} \bar{B}_{\frac{1}{2} I_4 s_4} \bar{\psi}_{a_2 \alpha_2 f_2} \psi_{a_2 \beta_1 f_1} \rangle^{(0)} \Gamma_{\alpha_1 \beta_1}^{e^1} \Gamma_{\alpha_2 \beta_2}^{-e^1}, \end{aligned} \quad (4.31)$$

with all fields at a same site and where the prime here means the sum over $I_1, \dots, I_4$, with $I_1 + I_2 = 0 = I_3 + I_4$. A similar expression holds for the components 12 ($s_1 s_2 t_3 t_4$), with $(1/2, I_3, s_3)$ and $(1/2, I_4, s_4)$, respectively, replaced by $(3/2, J_3, t_3)$ and $(3/2, J_4, t_4)$; with $(1/2, I_1, s_2)$ and $(1/2, I_2, s_2)$, respectively, replaced by $(3/2, J_1, t_1)$ and $(3/2, J_2, t_2)$ for the component 21 ($t_1 t_2 s_3 s_4$); and with $(1/2, I_1, s_1)$, $(1/2, I_2, s_2)$, $(1/2, I_3, s_3)$ and $(1/2, I_4, s_4)$, respectively, replaced by

$(3/2, J_1, t_1)$, $(3/2, J_2, t_2)$, $(3/2, J_3, t_3)$ and $(3/2, J_4, t_4)$ for the component 22 ($t_1 t_2 t_3 t_4$).

Taking into account the other directions and using translation and rotation invariance, we obtain for the kernel of $K^{(2)}$, in the total spin basis

$$\begin{aligned} K^{(2)}(x_1, x_2; x_3, x_4) &= \frac{1}{24} v^{(2)} \sum_{\sigma, j} \delta(x_2 - x_1 - \sigma e^j) \times \\ &\delta(x_1 - x_3) \delta(x_2 - x_4) \delta(x_2^0 - x_3^0), \end{aligned} \quad (4.32)$$

where here $v^{(2)}$ is the total spin basis matrix.

By a lengthy computation (see Appendix E), in the individual spin basis, the symmetric matrix $v^{(2)} = [v_{ij}]_{i, j=1, \dots, 20}$ has the following nonzero elements: $v_{11} = -1/3 = v_{44}$; $v_{17} = -4/\sqrt{6} = v_{113} = v_{412} = v_{418}$; $v_{110} = 4\sqrt{2}/3 = v_{415}$; $v_{22} = 10/3 = v_{33}$; $v_{23} = -11/3$; $v_{28} = 2\sqrt{2} = v_{317}$; $v_{211} = -4\sqrt{2}/3 = v_{314}$; $v_{311} = 2\sqrt{2}/3 = v_{214}$; $v_{55} = -3 = v_{2020}$; $v_{66} = -2 = v_{99} = v_{1616} = v_{1919}$; $v_{69} = -1 = v_{77} = v_{811} = v_{1212} = v_{1313} = v_{1417} = v_{1619} = v_{1818}$; $v_{710} = -2\sqrt{3} = v_{1013} = v_{1215} = v_{1518}$; $v_{1010} = -5/3 = v_{1515}$; $v_{1111} = -4/3 = v_{1114} = v_{1414}$.

It turns out that $v^{(2)}$ has an especially simple, almost diagonal form, in the total spin basis. Using the basis transformation $\mathcal{A}$ given in Section 4.1, it is written as

$$v^{(2)} = W \oplus W \oplus W \oplus W_{1d} \oplus W_2 \oplus W_{2d}, \quad (4.33)$$

with $W = \frac{1}{3} \begin{pmatrix} -1 & -\sqrt{80} \\ -\sqrt{80} & 1 \end{pmatrix}$ (with eigenvalues ± 3), $W_2 = \begin{pmatrix} 7 & 4 \\ 4 & 1 \end{pmatrix}$ (with eigenvalues -1 and 9), $W_{1d} = -3P_{13}$, $W_{2d} = -P_{22}$, where P_{13} is the orthogonal projection on the 7th through 13th components and P_{22} on the 16th through 20th.

The interaction present in Eq. (4.31) arises from a quark-antiquark exchange but, by inspection of the meson particle fields (see Refs. [6, 10]), this is *not* a meson particle exchange. To see this, consider the field averages in Eq. (4.31). The spin indices of the B and $\bar{B}$ fields are both lower. This forces the spin indices of the $\bar{\psi}\psi$ fields, in each of the two averages, to be either both lower or both upper. A true meson particle has one upper and one lower spin index. We refer to this interaction as a *quasi-meson exchange*.

Using the total spin basis representation for $v^{(2)}$, we return to the determination of bound states and solve the B-S equation using L (see Refs. [9, 10] for details). In terms of relative coordinates, we have the space range-one energy-independent potential

$$\hat{L}(\xi, \eta, k^0) = \kappa^2 v^{(2)} \sum_{j, \sigma} \delta(\vec{\eta} - \vec{\xi}) \delta(\sigma e^j - \vec{\xi}), \quad (4.34)$$

and the B-S equation, in the ladder approximation, has

the solution,

$$\begin{aligned} \hat{M}(\vec{\xi}, \vec{\eta}, k^0) &= \hat{M}_0(\vec{\xi}, \vec{\eta}) + \sum_{\sigma_1, \sigma_2, j_1, j_2} \hat{M}_0(\vec{\xi}, \sigma_1 \vec{e}^{j_1}) \times \\ &\quad \kappa^2 v^{(2)} \left(1 - \kappa^2 v^{(2)} \check{M}_0 \right)^{-1} (\sigma_1 \vec{e}^{j_1}, \sigma_2 \vec{e}^{j_2}) \times \\ &\quad \hat{M}_0(\sigma_2 \vec{e}^{j_2}, \eta), \end{aligned} \quad (4.35)$$

where the $\check{M}_0$ lattice indices are restricted to the spatial nearest neighbors of zero, and we suppress the k^0 dependence and spin indices on the r.h.s. Reinstating the spin indices, the resulting matrix is 120×120 , where we have implicit matrix multiplication in the spin indices $\check{M}_0$ means $\hat{M}_0$ with indices restricted by the multiplication of $v^{(2)}$ and we suppress k^0 dependence on the r.h.s.

To go further, we need properties of $\hat{M}_0$. In Appendix F, using the spectral representation for the two-baryon functions in Eq. (4.24), after passing to lattice relative coordinates, we derive a general spectral representation for $\hat{M}_0$. Assuming that the two-baryon function is diagonal in the spin indices in the individual spin basis [which holds up to and including $\mathcal{O}(\kappa^5)$, see Ref. [5], we show that the dominant contribution is given by, in the total spin basis,

$$\begin{aligned} \hat{M}_{0, SS_z S'_z}(\vec{\xi}, \vec{\eta}, k^0) &= \left[-2(2\pi)^{-3} \int_{\mathbb{T}^3} \tilde{G}(\vec{p}) \tilde{G}(\vec{p}) d\vec{p} \right. \\ &\quad \left. -2(2\pi)^3 \int_{-1}^1 \int_{-1}^1 \int_{\mathbb{T}^3} f(k^0, \lambda^0 \lambda'^0) d\lambda^0 \alpha_{\vec{p}}(\lambda^0) \right. \\ &\quad \left. \times d_{\lambda^0} \alpha_{\vec{p}}(\lambda^0) \Xi(\vec{p}, \vec{\xi}, \vec{\eta}) d\vec{p} \right] \delta_{SS'} \delta_{S_z S'_z}, \end{aligned} \quad (4.36)$$

where

$$\Xi(\vec{p}, \vec{\xi}, \vec{\eta}) = \delta_{S1} \cos \vec{p} \cdot \vec{\xi} \cos \vec{p} \cdot \vec{\eta} + \delta_{S0} \sin \vec{p} \cdot \vec{\xi} \sin \vec{p} \cdot \vec{\eta};$$

and the same for $T = 1, 3$ ($T = 0, 2$) replacing $S = 1$ ($S = 0$).

To continue, we set $k^0 = i(2\bar{m} - \epsilon)$, where we recall that $\bar{m}$ denotes the minimum of the baryon masses and $\epsilon > 0$ is the bound state binding energy. Recalling the definition of p_ℓ^2 given in Eq. (3.14), we further make the following approximations: a) we retain only the product of one-particle contributions; b) $w(\vec{p}) - m \approx \kappa^3 p_\ell^2 / 8$; c) $\bar{m} \approx -3 \ln \kappa$ and d) $Z(\vec{p}) \approx -(2\pi)^{-3} e^{-w(\vec{p})} \approx -(2\pi)^{-3} \kappa^3$. The approximate $\hat{M}_0$ becomes

$$\begin{aligned} \hat{M}_{0, SS_z S'_z} &= -\frac{2}{(2\pi)^3} \int_{\mathbb{T}^3} \left[1 + \frac{1}{e^{\kappa^3 p_\ell^2 / 4 + \epsilon} - 1} \right] \\ &\quad \times \Xi(\vec{p}, \vec{\xi}, \vec{\eta}) d\vec{p} \delta_{SS'} \delta_{S_z S'_z}. \end{aligned} \quad (4.37)$$

With all these approximations, the singularities of $\hat{M}$, below the two-baryon threshold $2\bar{m}$, occur as zeroes of the determinant of the 120×120 matrix $1 - \kappa^2 v^{(2)} \hat{M}_0(\sigma_1 \vec{e}^{j_1}, \sigma_2 \vec{e}^{j_2})$. Anticipating that the binding energy ϵ is of order κ^2 , we also approximate the denominator in $\hat{M}_0$ to get

$$\hat{M}_{0, SS_z S'_z}(\vec{\xi}, \vec{\eta}, k^0) \approx -\frac{2}{(2\pi)^3} (1 - e^{-\epsilon})^{-1} \int_{\mathbb{T}^3} \Xi(\vec{p}, \vec{\xi}, \vec{\eta}) d\vec{p},$$

and the same for T 's instead of S 's. These approximations are consistent with finding a bound state solution with binding energy greater than $\mathcal{O}(\kappa^3)$. Note that $\hat{M}_0(\sigma_1 \vec{e}^{j_1}, \sigma_2 \vec{e}^{j_2})$ is diagonal in j_1 and j_2 , and that the integral takes the values $\pm(2\pi)^3/2$. Thus, the bound state condition becomes

$$\det[1 + 4(1 - e^{-\epsilon})^{-1} v^{(2)} \kappa^2] = 0, \quad (4.38)$$

for a 20×20 matrix. Next, using the spectral representation for $v^{(2)}$, if λ is a negative eigenvalue, then the bound state condition is

$$1 + 4\lambda(1 - e^{-\epsilon})^{-1} \kappa^2 = 0,$$

with approximate solution $\epsilon = 4|\lambda|\kappa^2$, which is the bound state binding energy. Hence, we have a bound state for each negative eigenvalue of $v^{(2)}$, with multiplicity given by the space dimension 3. For these bound states we have attractive potential wells at $\pm e^j$, $j = 1, \dots, d = 3$, which would give us a multiplicity $2d$. However, we only have symmetric or antisymmetric wave functions, which reduces the multiplicity to $d = 3$.

Referring to the spectral decomposition of $v^{(2)}$ in the total spin basis, and recalling the ordering of the total spin basis, we see that the most strongly bound, bound states ($\lambda = -3$) are associated with a superposition of $p - n$ and $\Delta - \Delta$ total spin 1 states; and also the $\Delta - \Delta$ total spin 3 states. The more weakly bound, bound states ($\lambda = -1$) are associated with the superposition of $p - n$ and $\Delta - \Delta$ total spin 0 states; as well as the $\Delta - \Delta$ total spin 2 states.

We now explain the superposition of two-baryon states occurring in the bound state associated with the superposition of $p - n$ and $\Delta - \Delta$. Similar considerations apply to the other bound states. Going back to the decomposition of $v^{(2)}$ given in Eq. (4.33), this state is associated with the negative eigenvalue -3 of one of the three 2×2 W 's. Consider the first W and label the basis elements $(S, S_z) = (1, 1)$, $(T, T_z) = (1, 1)$ by $\sigma = 1, 2$. The corresponding eigenvector is $w_{-3} = (\sqrt{5}/3, 2/3)$. If we start from the B-S equation for (w_{-3}, Mw_{-3}) , it reduces to a single equation with $v^{(2)}$ substituted by -3 . Here, we have used the fact that $\hat{M}_0$ commutes with $v^{(2)}$ in our approximation. But (w_{-3}, Mw_{-3}) can be written, for $x_1^{(0)} = x_2^{(0)} \leq x_3^{(0)} = x_4^{(0)}$, as the average

$$\begin{aligned} &\langle [(\sqrt{5}/3) \mathcal{T}_{S=1, S_z=1}^{1/2}(x_1, x_2) + (2/3) \mathcal{T}_{T=1, T_z=1}^{3/2}(x_1, x_2)] \\ &\quad \times [(\sqrt{5}/3) \bar{\mathcal{T}}_{S=1, S_z=1}^{1/2}(x_3, x_4) + (2/3) \bar{\mathcal{T}}_{T=1, T_z=1}^{3/2}(x_3, x_4)] \rangle, \end{aligned}$$

so that the superposition of two-baryon states given by the field $[(\sqrt{5}/3) \bar{\mathcal{T}}_{S=1, S_z=1}^{1/2}(x_1, x_2) + (2/3) \bar{\mathcal{T}}_{T=1, T_z=1}^{3/2}(x_1, x_2)]$ more closely describes the field which creates the bound state.

4.4. Bound states for the $I = 3$ sector

In this subsection, we show that the application of our method to the maximum total isospin $I = 3$ sector (freezing all isospins to $+$) leads to *no* baryon-baryon bound states if $S = 2, 3$, but bound states do appear if $S = 0, 1$. The analysis here parallels that of the $I = 0$ case, but it is algebraically much simpler, and we will be brief.

The one-baryon states are associated with the fields $\bar{B}_t \equiv \bar{B}_{\frac{3}{2}t}$, $t = \pm 3/2, \pm 1/2$ of Eq. (3.9). The two-baryon states that we consider here, in the total $I = 3$, $I_z = 3$ sector, are those generated by the product of the above one-baryon fields $\bar{\tau}_{t_1 t_2}(x_1, x_2)$, $x_1^0 = x_2^0$. The bound states are formed with a quasi-meson exchange κ^2 space range-one potential which dominates and, in our ladder approximation, the space range-zero potential is zero.

After interchanging spin and isospin, the calculation of this potential is the *same* as for the lattice QCD model in $2 + 1$ dimensions and with 2×2 spin matrices and two flavors (see Ref. [9]), and the bound state spectral results are the same, apart from the additional space dimension.

In the total spin basis, the two-baryon states are given by

$$\mathcal{T}_{TT_z}(x_1, x_2) = \sum_{t_1+t_2=T_z} c_{t_1 t_2}^{TT_z} \tau_{t_1 t_2}(x_1, x_2),$$

where $c_{t_1 t_2}^{TT_z}$ are the C-G coefficients for the coupling of two spin $3/2$ states to give total spin $T = 3, 2, 1, 0$, and z -component T_z of total spin. They obey the symmetry property

$$\mathcal{T}_{TT_z}(x_1, x_2) = (-1)^T \mathcal{T}_{TT_z}(x_2, x_1),$$

for $T = 3, 1$; for $T = 0, 2$ ($1, 3$) they are space symmetric (antisymmetric). So, at coincident points, they vanish for $T = 3, 1$; for $T = 0, 2$ ($1, 3$) they are symmetric (anti-symmetric). Also, from the pointwise relations $\bar{B}_{\pm \frac{3}{2}} \bar{B}_{\pm \frac{3}{2}} = \bar{B}_{\pm \frac{3}{2}} \bar{B}_{\pm \frac{1}{2}} = \bar{B}_{\pm \frac{3}{2}} \bar{B}_{\mp \frac{1}{2}} = \bar{B}_{\pm \frac{1}{2}} \bar{B}_{\pm \frac{1}{2}} = 0$, we find that $\mathcal{T}_{T=2 T_z}$ is also zero at coincident points. A computation gives the coincident point linear dependency relation $\bar{B}_{\frac{3}{2}} \bar{B}_{-\frac{3}{2}} = -\bar{B}_{\frac{1}{2}} \bar{B}_{-\frac{1}{2}}$, and we find that $\tilde{\mathcal{T}}_{00} = 2\bar{B}_{\frac{3}{2}} \bar{B}_{-\frac{3}{2}}$.

The four-point correlation in the unmodified, shifted form, and in the individual spin basis, is taken as, with $x_1^0 = x_2^0$, $x_3^0 = x_4^0$, and for $x_1^0 \leq x_3^0$,

$$M_{t_1 t_2 t_3 t_4}(x_1, x_2, x_3, x_4) = \langle B_{t_1}(x_1) B_{t_2}(x_2) \bar{B}_{t_3}(x_3) \bar{B}_{t_4}(x_4) \rangle,$$

and with the usual interchange of barred and unbarred fields and taking the complex conjugate of the average, for $x_1^0 > x_3^0$. The associated uncorrelated or *Wickified* four baryon correlation is, for $x_1^0 \leq x_3^0$,

$$\begin{aligned} M_{0, t_1 t_2 t_3 t_4}(x_1, x_2, x_3, x_4) = & \\ & - \langle B_{t_1}(x_1) \bar{B}_{t_3}(x_3) \rangle \langle B_{t_2}(x_2) \bar{B}_{t_4}(x_4) \rangle \\ & + \langle B_{t_1}(x_1) \bar{B}_{t_4}(x_4) \rangle \langle B_{t_2}(x_2) \bar{B}_{t_3}(x_3) \rangle. \end{aligned}$$

The four-baryon correlations in the total spin basis are obtained by taking the corresponding C-G linear combinations. For $T = T' = T_z = T'_z = 0$, we obtain $M_{TT_z T' T'_z}^{(0)}(0) = -4$ and $M_{0, TT_z T' T'_z}^{(0)}(0) = -2$.

As in Section 4.3, an h -modified B-S equation and operator kernels are introduced. Then, the energy-independent space range-zero potential $K^{(0)} = [M_0^{(0)}]^{-1} - [M^{(0)}]^{-1} = 0$. Analogous to Eq. (4.30), the dominant interaction is a κ^2 quasi-meson space range-one potential which we take as our ladder approximation $L = K^{(2)} \kappa^2$ to K . Here, with $K^{(2)}$ and $v^{(2)}$ in the total spin basis, as in Eq. (4.32)

$$K^{(2)}(x_1, x_2; x_3, x_4) = \frac{1}{24} v^{(2)} \sum_{\sigma, j} \delta(x_2 - x_1 - \sigma e^j) \times \delta(x_1 - x_3) \delta(x_2 - x_4) \delta(x_2^0 - x_3^0),$$

where, in the individual spin basis,

$$\begin{aligned} v_{t_1, t_2, t_3, t_4}^{(2)} = & \Gamma_{\alpha_1 \beta_1}^{e_1} \Gamma_{\alpha_2 \beta_2}^{-e_1} \langle B_{t_1} \bar{B}_{t_3} \bar{\psi}_{a_1 \alpha_1 f_1} \psi_{a_1 \beta_2 f_2} \rangle^{(0)} \\ & \times \langle B_{t_2} \bar{B}_{t_4} \bar{\psi}_{a_2 \alpha_2 f_2} \psi_{a_2 \beta_1 f_1} \rangle^{(0)}. \end{aligned} \quad (4.39)$$

Next, using similar arguments to those in Appendix E, we can write

$$\begin{aligned} v_{t_1, t_2, t_3, t_4}^{(2)} = & \langle B_{t_1} \bar{B}_{t_3} : \bar{\psi}_{a_1 \alpha_1 f_1} \psi_{a_1 \beta_2 f_2} : \rangle^{(0)} \\ & \times \langle B_{t_2} \bar{B}_{t_4} : \bar{\psi}_{a_2 \beta_2 f_2} \psi_{a_2 \alpha_1 f_1} : \rangle^{(0)}, \end{aligned}$$

where in the expansion of the $\kappa = 0$ average using Wick's theorem the Wick ordering $::$ forbids the contraction between the *explicit internal* $\bar{\psi}$ and ψ fields. This restriction implies that $f_1 = f_2 = +$. In order to calculate $v_{t_1, t_2, t_3, t_4}^{(2)}$, we also take into account spin constraints and use conjugation and the $\kappa = 0$ spin flip symmetry.

Doing this and suppressing the $+$ isospin indices of the $\tilde{\psi}$, it is enough to calculate

$$\begin{aligned} \langle B_{\frac{1}{2}} \bar{B}_{\frac{1}{2}} : \bar{\psi}_+ \psi_+ : \rangle^{(0)} &= -2 = 2 \langle B_{\frac{1}{2}} \bar{B}_{\frac{1}{2}} : \bar{\psi}_- \psi_- : \rangle^{(0)}, \\ \langle B_{\frac{3}{2}} \bar{B}_{\frac{1}{2}} : \bar{\psi}_+ \psi_- : \rangle^{(0)} &= -\sqrt{3}, \\ \langle B_{\frac{3}{2}} \bar{B}_{\frac{3}{2}} : \bar{\psi}_+ \psi_+ : \rangle^{(0)} &= -3, \end{aligned}$$

and $\langle B_{\frac{1}{2}} \bar{B}_{-\frac{1}{2}} : \bar{\psi}_+ \psi_- : \rangle^{(0)} = -2$.

In this way, in the individual spin basis, using the same ordering as in the zero isospin case, for the 16×16 submatrix, the symmetric matrix $v^{(2)} = [v_{ij}]_{i,j=1,\dots,16}$ has the following nonzero elements: $v_{11} = 9 = v_{16 16}$; $v_{22} = 6 = v_{55} = v_{12 12} = v_{15 15}$; $v_{25} = 3 = v_{33} = v_{47} = v_{88} = v_{99} = v_{10 13} = v_{12 15} = v_{14 14}$; $v_{36} = 2\sqrt{3} = v_{69} = v_{8 11} = v_{11 14}$; $v_{66} = 5 = v_{11 11}$; $v_{77} = 4 = v_{7 10} = v_{10 10}$. The matrix is easily diagonalized in the following re-ordered basis: 3, 6, 9, 4, 7, 10, 13, 8, 11, 14, 2, 5, 12, 15, 1, 16. Letting $\mathcal{V}$ denote this matrix, we have $\mathcal{V} = \mathcal{V}_1 \oplus \mathcal{V}_2 \oplus \mathcal{V}_1 \oplus \mathcal{V}_3 \oplus \mathcal{V}_3 \oplus \mathcal{V}_4 \oplus \mathcal{V}_4$, with $\mathcal{V}_{1,2,3,4}$ symmetric, and where $\mathcal{V}_1$ is 3×3 with nonvanishing elements $\mathcal{V}_{1;11} = 3 = \mathcal{V}_{1;33}$, $\mathcal{V}_{1;22} = 5$, $\mathcal{V}_{1;12} = 2\sqrt{3} = \mathcal{V}_{1;23}$; $\mathcal{V}_2$ is 4×4 with nonvanishing elements $\mathcal{V}_{2;22} = 4 = \mathcal{V}_{2;33} = \mathcal{V}_{2;23}$, $\mathcal{V}_{2;12} = 3 = \mathcal{V}_{2;34}$; $\mathcal{V}_3$ is 2×2 with nonvanishing

elements $\mathcal{V}_{3;11} = 6 = \mathcal{V}_{3;22}$, $\mathcal{V}_{3;12} = 3$; $\mathcal{V}_4$ is 1×1 with nonvanishing elements $\mathcal{V}_{4;11} = 9$.

Concerning their eigenvalues and eigenvectors (T, T_z) (the total and z -component of the total spin), we have: 9 (3, 1), 3 (2, 1) and -1 (1, 1) for $\mathcal{V}_1$ in the first spot; 9 (3, 0), 3 (2, 0), -1 (1, 0) and -3 (0, 0) for $\mathcal{V}_2$; 9 (3, -1), 3 (2, -1) and -1 (1, -1) for $\mathcal{V}_1$ in the second spot; 9 (3, 2) and 3 (2, 2) for $\mathcal{V}_3$ first spot; 9 (3, -2) and 3 (2, -2) for $\mathcal{V}_3$ in the second spot; 9 (3, 3) for $\mathcal{V}_4$ in the first spot; 9 (3, -3) for $\mathcal{V}_4$ in the second spot.

As we see, in the total spin basis, $v^{(2)}$ is diagonal. Alternatively, it can be also diagonalized by the orthogonal transformation relating the total and individual spin basis. Similar to what occurs for the $I = 0$ sector, we shall see that the negative eigenvalues are associated with bound states. Referring to the above, the negative eigenvalue -3 (-1) is associated with $T, T_z = (0, 0)$ ($T, T_z = (1, 1), (1, 0), (1, -1)$).

Proceeding as in the $I = 0$ case, in terms of lattice relative coordinates, the space range-one energy-independent potential in the ladder approximation is, as in Eq. (4.34), given by

$$\hat{L}(\vec{\xi}, \vec{\eta}, k^0) = \kappa^2 v^{(2)} \sum_{j, \sigma} \delta(\vec{\eta} - \vec{\xi}) \delta(\sigma e^j - \vec{\xi}),$$

with $v^{(2)}$ in the total spin basis and the B-S equation has the solution given by Eq. (4.35). Here $\hat{M}_0(\vec{\xi}, \vec{\eta}, k^0)$ is derived as in Appendix F and in the total spin representation is given as in Eq. (4.36) with S, S_z replaced by T, T_z , as well as their primes and with now

$$\begin{aligned} \Xi(\vec{p}, \vec{\xi}, \vec{\eta}) &= (\delta_{T2} + \delta_{T0}) \cos \vec{p} \cdot \vec{\xi} \cos \vec{p} \cdot \vec{\eta} \\ &+ (\delta_{T3} + \delta_{T1}) \sin \vec{p} \cdot \vec{\xi} \sin \vec{p} \cdot \vec{\eta}. \end{aligned}$$

Finally, approximating $\hat{M}_0$ as before, for each negative eigenvalue λ of $v^{(2)}$ we have the bound state condition $1 + 4\lambda(1 - e^{-\epsilon})^{-1}\kappa^2 = 0$, with the approximate solution of binding energy $\epsilon = 4|\lambda|\kappa^2$. Hence, we only have bound states of spin one and zero, with multiplicities 3 and 1, respectively.

5. CONNECTION BETWEEN THE B-S EQUATION AND A LATTICE SCHRÖDINGER EQUATION

In this section, to understand our spectral results in a simple way, we exploit the similarity between our lattice B-S equation for $\hat{M}(\vec{\xi}, \vec{\eta}, k^0)$ and a one-particle Schrödinger resolvent operator equation for the lattice Hamiltonian

$$H = H_0 + V,$$

with a kinetic energy operator H_0 and a potential energy operator V .

For κ and ϵ small, we further approximate $\hat{M}_0(\vec{\xi}, \vec{\eta}, k^0)$ of Eq. (4.37) by

$$\hat{M}_0 = -\frac{2}{(2\pi)^3} \int_{\mathbb{T}^3} \frac{1}{\kappa^3 p_\ell^2/4 + \epsilon} \Xi(\vec{p}, \vec{\xi}, \vec{\eta}) d\vec{p},$$

for p_ℓ given in Eq. (3.14) and Ξ depends on the total spin.

Consider the Schrödinger resolvent equation,

$$(H - z)^{-1} = (H_0 - z)^{-1} - (H_0 - z)^{-1} V (H - z)^{-1},$$

for z not in the spectrum of H_0 and H .

Upon comparing with minus the B-S equation in our ladder approximation, i.e. $-\hat{M} = -\hat{M}_0 - \hat{M}_0 \hat{L} \hat{M}$, we can make the identifications

$$H_0 = \frac{\kappa^3}{8} (-\Delta_\ell) \quad , \quad \hat{L} = V \quad , \quad z = -\frac{\epsilon}{2} \quad (z < 0),$$

where Δ_ℓ is the lattice Laplacian with Fourier transform $-p_\ell^2$. The Hamiltonian $H = H_0 + V$ acts on the space $\ell_{20}^2(\mathbb{Z}^3)$, which decomposes into a direct sum of space-symmetric and antisymmetric subspaces depending on the total spin indices (omitted here for simplicity of notation!). For $f \in \ell_{20}^2(\mathbb{Z}^3)$, for each x , $f_R(x)$ is the R th component where R runs over the 20 total spin basis indices. The subspace is symmetric (antisymmetric) for $\Xi(\vec{p}, \vec{\xi}, \vec{\eta}) = \cos \vec{p} \cdot \vec{\xi} \cos \vec{p} \cdot \vec{\eta}$ ($\sin \vec{p} \cdot \vec{\xi} \sin \vec{p} \cdot \vec{\eta}$), corresponding to total spin $S = 1, T = 1, 3$ ($S = 0, T = 0, 2$).

The kernel of V is

$$V(\vec{\xi}, \vec{\eta}) = \kappa^2 v^{(2)} \sum_{\sigma, j} \delta(\vec{\xi} - \sigma e^j) \delta(\vec{\xi} - \vec{\eta}),$$

and, for small κ , V dominates H_0 .

In this case, the negative energy bound state eigenfunctions of H with eigenvalue $\lambda = z < 0$ are those of V which are easily seen to have components

$$f_R^{(k)}(x) = \alpha_R [\delta(x - e^k) \pm \delta(x + e^k)],$$

where α is an eigenvector of the eigenvalue equation

$$\kappa^2 v^{(2)} \alpha = \lambda \alpha.$$

The plus (minus) sign occurs for the symmetric (antisymmetric) subspace of $\ell_{20}^2(\mathbb{Z}^3)$. The eigenvectors are localized at $\pm e^k$, respectively. When the free Hamiltonian H_0 is not neglected, the negative eigenvectors of H get spread out.

We now reinstate H_0 and obtain the spectral representation of H where we write $V(\vec{\xi}, \vec{\eta}) = 2\kappa^2 \mu \sum_j \delta(\vec{\xi} - e^j) \delta(\vec{\xi} - \vec{\eta})$, where μ is a negative eigenvalue of $v^{(2)}$, and we are in the space symmetric or antisymmetric subspace. As H acts in the space of symmetric or antisymmetric subspace of $\ell_{20}^2(\mathbb{Z}^3)$ the $\delta(\vec{\xi} + e^j)$ term in the potential can be dropped and the term $\delta(\vec{\xi} - e^j)$ gets multiplied by 2. The space symmetry or antisymmetry is incorporated

automatically in the definition of $(H_0 - z)^{-1}$ through the function $\Xi(\vec{p}, \vec{\xi}, \vec{\eta})$.

The spectral results below hold for all $\kappa > 0$ while the correspondence with the B-S equation is valid only for $\kappa \ll 1$. With the above observations, the resolvent equation becomes

$$(H - z)^{-1}(\vec{\xi}, \vec{\eta}) = (H_0 - z)^{-1}(\vec{\xi}, \vec{\eta}) - (H_0 - z)^{-1}(\vec{\xi}, e^j) \times 2\kappa^2\mu(H - z)^{-1}(e^j, \vec{\eta}),$$

with solution

$$(H - z)^{-1}(\vec{\xi}, \vec{\eta}) = (H_0 - z)^{-1}(\vec{\xi}, \vec{\eta}) - (H_0 - z)^{-1}(\vec{\xi}, e^j) \times 2\kappa^2\mu B_{jk}(H_0 - z)^{-1}(e^k, \vec{\eta}),$$

where $B \equiv A^{-1}$ and $A = [1 + (H_0 - z)^{-1} 2\kappa^2\mu]$ restricted to $\ell_2(\{e^1, e^2, e^3\})$ so it is a 3×3 matrix. The matrix A has the structure

$$A = \begin{pmatrix} a & b & b \\ b & a & b \\ b & b & a \end{pmatrix},$$

where $a = 1 + 2\kappa^2\mu(H_0 - z)^{-1}(\vec{r}, \vec{t})$, $(\vec{r}, \vec{t}) = (e^1, e^1)$, (e^2, e^2) or (e^3, e^3) , and $b = 2\kappa^2\mu(H_0 - z)^{-1}(\vec{r}, \vec{t})$, $(\vec{r}, \vec{t}) = (e^1, e^2)$, (e^1, e^3) or (e^2, e^3) . Furthermore, $\det A = (a - b)^2(a + 2b)$, and the eigenvalues of A are $\lambda_1 = \lambda_2 = (a - b)$, $\lambda_3 = a + 2b$, with corresponding normalized eigenvectors $f_1 = (r_1, r_2, r_3)$, $f_2 = (t_1, t_2, t_3)$, $f_3 = \frac{1}{\sqrt{3}}(1, 1, 1)$, where $\sum_i r_i = \sum_i t_i = 0$ and $f_1 \perp f_2$.

The matrix B has the same structure as A and is given by

$$B = \frac{1}{(a - b)(a + 2b)} \begin{pmatrix} a + b & -b & -b \\ -b & a + b & -b \\ -b & -b & a + b \end{pmatrix}.$$

The negative energy bound states occur as zeroes of the denominator of B , and are given by the zeroes of $(a - b)$ and $(a + 2b)$.

Denoting minus the binding energies by $z = -\epsilon$, for $\epsilon_1 = \epsilon_2, \epsilon_3$, we have the conditions, for $z = -\epsilon_1 = -\epsilon_2$

$$0 = 1 + 2\kappa^2\mu[(H_0 - z)^{-1}(e^1, e^1) - (H_0 - z)^{-1}(e^1, e^2)],$$

and, for $z = -\epsilon_3$

$$0 = 1 + 2\kappa^2\mu[(H_0 - z)^{-1}(e^1, e^1) + 2(H_0 - z)^{-1}(e^1, e^2)].$$

The corresponding numerators of the resolvents $(H_0 - z)^{-1}$, in Fourier space, are $(\cos p^1)^2 - \cos p^1 \cos p^2$ and $(\cos p^1)^2 + 2 \cos p^1 \cos p^2$, in addition to the factor $\Xi(\vec{p}, \vec{\xi}, \vec{\eta})$. Expanding the denominator $(p_\ell^2 - z)$ shows that $\epsilon_3 > \epsilon_1$.

Recalling that the resolvent contains all spectral information, we now obtain the bound state eigenfunctions. To this end, from the spectral theorem

$$\begin{aligned} \lim_{z \rightarrow -\epsilon_i} (-\epsilon_i - z)(H - z)^{-1}(\vec{\xi}, \vec{\eta}) &= \sum_{j=1}^{m_i} \psi_{ij}(\vec{\xi}) \bar{\psi}_{ij}(\vec{\eta}) \\ &= -(H_0 + \epsilon_i)^{-1}(\vec{\xi}, e^j) F_{ijk}(H_0 + \epsilon_i)^{-1}(e^k, \vec{\eta}), \end{aligned}$$

where $F_{ijk} = \lim_{z \rightarrow -\epsilon_i} 2\kappa^2\mu B_{jk}$ and $\psi_{ij}(\vec{\xi})$ are eigenfunctions of H with eigenvalue $-\epsilon_i$; $j = 1, 2, \dots, m_i$ labels the m_i degenerate states.

For $i = 3$, $a = -2b$ and

$$F_3 = \frac{1}{3c_3} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix},$$

where, upon expanding $(a + 2b)$ in the denominator of B in powers of $(z + \epsilon_3)$, we have $c_3 = 2[(H_0 - z)^{-2}(e^1, e^1) + 2(H_0 - z)^{-1}(e^1, e^2)]|_{z=-\epsilon_3}$. The eigenvalues of the matrix are 0 and 3, with multiplicity two and one, respectively. The corresponding eigenvectors are h_1, h_2 and $h_3 = (1, 1, 1)/\sqrt{3}$; in h_1 and h_2 the sum of the components is zero and h_1 is orthogonal to h_2 . Noting that $h_{3j}h_{3k}$ is the jk matrix element of the orthogonal projection on the multiplicity one eigenspace, we have

$$\psi_3(\vec{\xi})\psi_3(\vec{\eta}) = \frac{1}{c_3}(H_0 + \epsilon_1)^{-1}(\vec{\xi}, e^j)h_{3j}h_{3k}(H_0 + \epsilon_1)^{-1}(e^k, \vec{\eta}),$$

so that the normalized eigenfunction is

$$\psi_3(\vec{\xi}) = \frac{1}{\sqrt{c_3}}(H_0 + \epsilon_1)^{-1}(\vec{\xi}, e^j)h_{3j}.$$

Similarly, for the doubly degenerate eigenvalue $\epsilon_1 = \epsilon_2$, we have

$$F_1 = \frac{1}{3c_1} \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{pmatrix},$$

and the matrix has eigenvalues 0 (multiplicity one) and 3 (multiplicity two) and eigenvectors $h_3 = (1, 1, 1)/\sqrt{3}$; and h_1, h_2 where h_1 is orthogonal to h_2 and the sum of components is zero. Hence,

$$\begin{aligned} \sum_{m=1,2} \psi_{1m}(\vec{\xi})\psi_{1m}(\vec{\eta}) &= \frac{1}{c_1}(H_0 + \epsilon_1)^{-1}(\vec{\xi}, e^j)[h_{1j}h_{1k} \\ &\quad + h_{2j}h_{2k}](H_0 + \epsilon_1)^{-1}(e^k, \vec{\eta}), \end{aligned}$$

so that the normalized eigenfunctions ψ_{11} and ψ_{12} are given by

$$\psi_{11}(\vec{\xi}) = \frac{1}{\sqrt{c_1}}(H_0 + \epsilon_1)^{-1}(\vec{\xi}, e^j)h_{1j},$$

and

$$\psi_{12}(\vec{\xi}) = \frac{1}{\sqrt{c_1}}(H_0 + \epsilon_1)^{-1}(\vec{\xi}, e^j)h_{2j},$$

which are orthogonal to each other since

$$\sum_{\vec{\xi}} \psi_{11}(\vec{\xi})\psi_{12}(\vec{\xi}) = \frac{1}{c_1}h_{1j}(H_0 + \epsilon_1)^{-2}(e^j, e^k)h_{2k},$$

and from the structure of $(H_0 + \epsilon_1)^{-1}$, h_1 and h_2 are also its orthogonal eigenvectors.

The eigenfunctions $\psi_{11,12,3}(\vec{\xi})$ have exponential decay by the Paley-Wiener theorem (see Ref. [36]), and in fact the integrals of the form $(H_0 + \epsilon)^{-1}(\vec{\xi}, \vec{\eta})$ occurring in the wave functions as well as in the bound state condition can be evaluated explicitly in terms of elliptic functions (see Refs. [37, 38, 39, 40, 41, 42] and also Refs. [43, 44]). We remark that another approach to the eigenfunctions is group theoretical noting that the potential is invariant under the symmetric group S_3 of permutations of the coordinate components. We see that the eigenfunction ψ_3 is a basis for the one-dimensional identity representation of S_3 ; ψ_{11} , ψ_{12} form a basis for the two-dimensional representation of S_3 .

6. CONCLUSIONS AND FINAL REMARKS

We point out that we can determine the baryon-baryon bound state spectrum using the more traditional unmodified B-S equation (4.23). The solution of the B-S equation is more complicated as it deals with a three-part energy-dependent potential, as explained in Appendix D. Using this approach, we do obtain the same bound state results.

To conclude, if we consider contributions to the B-S kernel comprised of linear chains of quark, anti-quark pairs they result in an exponentially decreasing potential with decay rate $-2 \ln \kappa$, as for the Yukawa theory. Surely, it would be interesting to look for the expected *distance*⁻¹ Ornstein-Zernicke like correction to this potential and to determine the spin and isospin dependencies of the binding energy, and the dependence on the number of flavors. Especially, the analysis of the SU(3) flavor case can shed some light in understanding bound states when strangeness and, consequently, Λ particles are present. Also, there is the problem of determining bound states of more than two baryons. However, lattice effects are expected to be relevant and unrealistic in determining the resulting geometric spatial configuration of possible bound states.

We make some remarks on pitfalls that can occur when trying to determine bound states by simulation methods. For example, consider the total isospin $I = 3$, $I_z = 3$ sector treated in Section 4.4. We have found bound states with binding energies proportional to $3\kappa^2$ and κ^2 which are associated with the total spin $(T, T_z) = (0, 0)$ and $(T, T_z) = (1, (-1, 0, 1))$, respectively. If we look at the correlation $\langle \mathcal{T}_{TT_z}(x_1, x_2) \bar{\mathcal{T}}_{TT_z}(x_3, x_4) \rangle$ with $x_1^0 < x_3^0$, $\vec{x}_1 = \vec{x}_2$ and $\vec{x}_3 = \vec{x}_4$, it is zero for $(T, T_z) = (1, (-1, 0, 1))$, as $\mathcal{T}_{TT_z}(x_1, x_2)$ is space anti-symmetric. Thus, these bound states would be missed! Even if we take $\vec{x}_1 \neq \vec{x}_2$ and $\vec{x}_3 \neq \vec{x}_4$, the decay rate in $x_3^0 - x_1^0$ would not be $2\bar{m} - \alpha\kappa^2$, $\alpha > 0$, but the smaller one of $2\bar{m} - 3\alpha\kappa^2$ since we expect there is overlap with the $(T, T_z) = (0, 0)$ state. This overlap occurs as there is *no* spin invariance and associated decomposition into disjoint spin sectors, at $\kappa \neq 0$. Thus it seems to be very difficult to see all bound states with small spectral separa-

rations.

Finally, and more importantly, we would like to know how the baryon spectrum and the bound state binding energies behave near the scaling limit.

This work was supported by CNPq and FAPESP. Algebraic computations were checked with Maple and Matlab, for which we thank Louis O'Carroll. We also thank P.H.R. dos Anjos for help in unveiling the time reflection symmetry of Appendix B.

APPENDIX A: Isospin orthogonality relations and correlation identities

Here we obtain isospin orthogonality relations for the two and four-baryon correlations used throughout this paper, and which hold for all κ . These relations also hold for spin at $\kappa = 0$, since the action has a SU(2) symmetry in the lower components of the spin index, and spin is also conserved under this condition.

We consider, typically, the two-point correlation $\langle \phi_{I_1 m_1}(x_1) \bar{\phi}_{I_2 m_2}(x_2) \rangle$ and the four-point correlation $\langle \phi_{I_1 m_1}(x_1) \phi_{I_2 m_2}(x_2) \bar{\phi}_{I_3 m_3}(x_3) \bar{\phi}_{I_4 m_4}(x_4) \rangle$, where we suppress the spin index and display only the I and I_z labels.

Considering first the two-point correlation and suppressing the gauge and spin indices and site arguments, the two-point correlation is given by linear combinations in the gauge and spin indices, and linear C-G combinations in the isospin indices of the basic correlation

$$\langle \psi_{i_1} \psi_{i_2} \psi_{i_3} \bar{\psi}_{i_4} \bar{\psi}_{i_5} \bar{\psi}_{i_6} \rangle \equiv T_{123;456}.$$

We start from the identity, with arbitrary $U \in U(2)$,

$$\begin{aligned} \langle \psi_{i_1} \psi_{i_2} \psi_{i_3} (\bar{\psi} U)_{i_4} (\bar{\psi} U)_{i_5} (\bar{\psi} U)_{i_6} \rangle \\ = \langle (U \psi)_{i_1} (U \psi)_{i_2} (U \psi)_{i_3} \bar{\psi}_{i_4} \bar{\psi}_{i_5} \bar{\psi}_{i_6} \rangle, \end{aligned}$$

where the r.h.s. is obtained from the l.h.s. using the two-flavor isospin symmetry. We schematically rewrite this relation as

$$(TU_3)_{123;456} = (U_3 T)_{123;456}, \quad (\text{A.1})$$

where we set $U_3 \equiv U \otimes U \otimes U$.

We obtain orthogonality relations for the two-baryon correlation by relating this identity to the usual quantum mechanical sum of angular momentum relations. We write $U = e^{i\theta\sigma/2}$, $I = \sigma/2$, where $\theta\sigma = \theta_x\sigma_x + \theta_y\sigma_y + \theta_z\sigma_z$, or $\theta_{x,y,z} \in \mathbb{R}$. Expanding in θ , we have $U = 1 + i\theta I - \frac{\theta^2}{2} + \dots$ such that

$$\begin{aligned} U_3 = & 1 \otimes 1 \otimes 1 + i\theta(I \otimes 1 \otimes 1 + 1 \otimes I \otimes 1 + 1 \otimes 1 \otimes I) \\ & - \frac{\theta^2}{2}(I^2 \otimes 1 \otimes 1 + 1 \otimes I^2 \otimes 1 + 1 \otimes 1 \otimes I^2 \\ & + 2I \otimes I \otimes 1 + 2I \otimes 1 \otimes I + 21 \otimes I \otimes I) \dots \end{aligned} \quad (\text{A.2})$$

Multiplying Eq. (A.1) by $\bar{w}_{123}$ on the left and v_{123} on the right, and equating the coefficients of $i\theta_z$ and $-(\theta_x^2 + \theta_y^2 + \theta_z^2)/2$, we obtain the identities

$$\bar{w}_{123}(TI_z)_{123;456}v_{456} = \bar{w}_{123}(I_z T)_{123;456}v_{456}, \quad (\text{A.3})$$

and

$$\bar{w}_{123}(TI^2)_{123;456}v_{456} = \bar{w}_{123}(I^2T)_{123;456}v_{456},$$

where, with an abuse of notation, $I_z = I_z \otimes 1 \otimes 1 + 1 \otimes I_z \otimes 1 + 1 \otimes 1 \otimes I_z$ and $I^2 = (\vec{I}_1 + \vec{I}_2 + \vec{I}_3)^2$, and $\vec{I}_j$, as usual, acts on the j th spot of the tensor product.

We will see that $I_z (I^2)$ has the interpretation of the z -component (square) of total isospin. We can use the usual complex Hilbert space notation $(\ , \)$ (as $\sigma_{x,y,z}$ are self-adjoint) and write the above as

$$(w, TI_z v) = (I_z w, T v), \quad (w, TI^2 v) = (I^2 w, T v). \quad (\text{A.4})$$

Thus, if v and w correspond to eigenfunctions with distinct eigenvalues we have orthogonality. To be more explicit, using the usual quantum mechanics notation for α (β) for spin up (down) state, and suppressing the tensor product notation, an isospin state of total isospin $\frac{3}{2}$ and z -component $\frac{3}{2}$ ($\frac{1}{2}$) are represented as the C-G linear combination $v_{\frac{3}{2}\frac{3}{2}} = \alpha\alpha\alpha$ and $v_{\frac{3}{2}\frac{1}{2}} = \frac{1}{\sqrt{3}}(\alpha\alpha\beta + \alpha\beta\alpha + \beta\alpha\alpha)$, respectively.

Taking these states into Eqs. (A.4), and using the usual quantum mechanical relations gives the orthogonality relations. Furthermore, taking the appropriate gauge and spin linear combinations to obtain $\phi_{I\ell m_\ell}$ gives the orthogonality relations for the two-baryon function.

Similar to the I_z identity of Eq. (A.3), we also have, with $I_\pm \equiv I_x \pm iI_y$,

$$(w, TI_\pm v) = (I_\pm w, T v). \quad (\text{A.5})$$

Next, we take $v = \chi_{Im}$ and $v = \chi_{I(m-1)}$, where $\chi_{I\ell}$ is the usual normalized eigenfunction of total isospin I and z -component ℓ . Hence, $\chi_{I\ell}$ satisfies $I_\pm \chi_{I\ell} = c_\ell^\pm \chi_{I(\ell \pm 1)}$, where

$$c_\ell^\pm = [I(I+1) - \ell(\ell \pm 1)]^{1/2}.$$

Substituting in Eq. (A.5), with the lower sign, and noting that $c_\ell^- = c_{\ell-1}^+$, gives

$$(\chi_{I(\ell-1)}, T \chi_{I(\ell-1)}) = (\chi_{I\ell}, T \chi_{I\ell}),$$

which implies the identity

$$\langle \phi_{I(\ell-1)}(x) \bar{\phi}_{I(\ell-1)}(y) \rangle = \langle \phi_{I\ell}(x) \bar{\phi}_{I\ell}(y) \rangle. \quad (\text{A.6})$$

Of course, Eq. (A.6) implies that all associated one-baryon spectral properties are the same, for I fixed, and all I_z .

Let us now turn to the four-baryon functions. The basic four-point function is

$$\langle \psi_{i_1} \dots \psi_{i_6} \bar{\psi}_{i_7} \dots \bar{\psi}_{i_{12}} \rangle \equiv T_{1\dots 6;7\dots 12}.$$

As before, we have the identity

$$TU_6 = U_6 T, \quad (\text{A.7})$$

where $U_6 \equiv \bigotimes_{j=1}^6 U$. Decomposing U_6 as $U_3 \otimes U_3$ and using similar considerations as above, we obtain orthogonality relations for the z -component and square of the total isospin by the appropriate $C - G$ linear combinations of the product of the one-particle isospin states $\phi_{I\ell}$.

Concerning the *Wickified* four baryon function M_0 of Section 4.3, the basic correlation is

$$\begin{aligned} T_{0,1\dots 6;7\dots 12} \equiv & -\langle \psi_1 \dots \psi_3 \bar{\psi}_7 \dots \bar{\psi}_9 \rangle \langle \psi_4 \dots \psi_6 \bar{\psi}_{10} \dots \bar{\psi}_{12} \rangle \\ & + \langle \psi_1 \dots \psi_3 \bar{\psi}_{10} \dots \bar{\psi}_{12} \rangle \langle \psi_4 \dots \psi_6 \bar{\psi}_7 \dots \bar{\psi}_9 \rangle. \end{aligned}$$

We have the identity

$$\begin{aligned} & -\langle \psi_1 \dots \psi_3 (\bar{\psi}U)_7 \dots (\bar{\psi}U)_9 \rangle \langle \psi_4 \dots \psi_6 \\ & \times (\bar{\psi}U)_{10} \dots (\bar{\psi}U)_{12} \rangle + \langle \psi_1 \dots \psi_3 (\bar{\psi}U)_{10} \dots (\bar{\psi}U)_{12} \rangle \\ & \times \langle \psi_4 \dots \psi_6 (\bar{\psi}U)_7 \dots (\bar{\psi}U)_9 \rangle = -\langle (U\psi)_1 \dots (U\psi)_3 \\ & \times \bar{\psi}_7 \dots \bar{\psi}_9 \rangle \langle (U\psi)_4 \dots (U\psi)_6 \bar{\psi}_{10} \dots \bar{\psi}_{12} \rangle \\ & + \langle (U\psi)_1 \dots (U\psi)_3 \bar{\psi}_{10} \dots \bar{\psi}_{12} \rangle \langle (U\psi)_4 \dots (U\psi)_6 \\ & \times \bar{\psi}_7 \dots \bar{\psi}_9 \rangle, \end{aligned}$$

which can be compactly written as

$$T_0 U_6 = U_6 T_0. \quad (\text{A.8})$$

We conclude by remarking that the transformation properties for T and T_0 of Eqs (A.7) and (A.8) are the same. Consequently, we have the same orthogonality relations for the z -component and square of total isospin of two-baryon states.

APPENDIX B: The hyperplane decoupling method

In this Appendix, we use the hyperplane decoupling method to see how the twenty baryon particles arise in the energy-momentum spectrum for our SU(3), two-flavor lattice QCD model. It is understood here that we first work in a finite-volume model and then with bounds on the normalized correlations which are uniform in the volume, obtained via a polymer expansion, the arguments can be extended to the infinite volume theory in a standard way (see Ref. [13, 33, 45]). This step is *not* treated here, and we work directly in the infinite volume. Also, we explicitly discuss the hyperplane decoupling method applied to the time direction, the treatment for the three space directions being similar.

In order to give a brief description of the method, we recall that the starting point for the time hyperplane decoupling method is to consider the model with interpolating parameters replacing the hopping terms giving couplings in the temporal direction. In terms of the model action, for the hopping terms connecting the hyperplanes $x^0 = p$ and $x^0 = p + 1$, the hopping parameter κ is replaced by a real parameter κ_p . (Note that setting $\kappa_p = 0$ erases all connections in the time direction between p and $p + 1$!) (see Ref. [46]). Doing so, as in Sections 3 and 4, we define κ_p -dependent correlations e.g. the κ_p dependent zero, two and four-baryon correlations, which provide a continuation to complex κ_p afterwards.

The main point here is that decay properties of these correlations in direct lattice spacetime are related to the vanishing of the first n derivatives with respect to κ_p , at $\kappa_p = 0$. In our case, the first two temporal hyperplane derivatives of G vanish, which leads to a $-3 \ln \kappa$ decay rate. The third derivative has an important product structure which implies that the first four derivatives of the G convolution inverse Γ are zero. This in turn shows that Γ has a faster decay rate of at least $-4 \ln \kappa$. Associated to these properties, by Fourier transform, there are analyticity properties in energy-momentum Fourier dual variables. The Fourier transform $\tilde{\Gamma}(p)$ is analytic in a larger p^0 strip than $\tilde{G}(p)$ so that $\tilde{\Gamma}^{-1}(p) = \text{cof}^t[\tilde{\Gamma}](p)/\det \tilde{\Gamma}(p)$ provides a meromorphic extension of $\tilde{G}(p)$ to the larger strip. As seen in Section 3, the dispersion curves are determined as the zeroes of $\det \tilde{\Gamma}(p^0 = iw(\vec{p}), \vec{p})$, so that they are isolated. This is a brief account of how, using the spectral representations for the Fourier transformed correlations, the method leads to spectral properties.

Next, we show how the 20 baryon particles arise in the two-flavor case, using the hyperplane decoupling method. Also, we show how to classify these states according to total isospin I , z -component isospin I_z and spin s_z . Let $\epsilon \psi_{\vec{\alpha}\vec{f}}^3 \equiv \epsilon_{abc} \psi_{a\alpha_1 f_1} \psi_{b\alpha_2 f_2} \psi_{c\alpha_3 f_3}$, which is invariant under the exchange $\alpha_j f_j \leftrightarrow f_k \alpha_k$, which we call the *totally symmetric property*. As seen below, we can restrict our analysis to the lower spin components, $\alpha_j = 3, 4$.

For $x^0 < y^0$, we consider a generic two-baryon correlation

$$G_{\vec{\alpha}_1 \vec{f}_1 \vec{\alpha}_2 \vec{f}_2}(x, y) \equiv \langle \epsilon \psi_{\vec{\alpha}_1 \vec{f}_1}^3(x) \epsilon \bar{\psi}_{\vec{\alpha}_2 \vec{f}_2}^3(y) \rangle.$$

We now consider the derivatives with respect to the hyperplane parameters κ_p , at $\kappa_p = 0$. It is easy to see that $G_{\vec{\alpha}_1 \vec{f}_1 \vec{\alpha}_2 \vec{f}_2}^{(0),(1),(2)} = 0$ because of imbalance of fermion fields ψ . For the third derivative, after performing the gauge integral using [see Ref. [2]]

$$\begin{aligned} \mathcal{I}_3 &\equiv \int U(g)_{a_1 b_1} U(g)_{a_2 b_2} U(g)_{a_3 b_3} d\mu(g) \\ &\equiv \int g_{a_1 b_1} g_{a_2 b_2} g_{a_3 b_3} d\mu(g) \\ &= \frac{1}{6} \epsilon_{a_1 a_2 a_3} \epsilon_{b_1 b_2 b_3}, \end{aligned} \quad (\text{B.1})$$

we get, with $\vec{w} \in \mathbb{Z}^3$,

$$\begin{aligned} G_{\vec{\alpha}_1 \vec{f}_1 \vec{\alpha}_2 \vec{f}_2}^{(3)}(x, y) &= -\frac{1}{36} \sum_{\vec{w}, \vec{\beta}, \vec{h}} \langle \epsilon \psi_{\vec{\alpha}_1 \vec{f}_1}^3(x) \epsilon \bar{\psi}_{\vec{\beta} \vec{h}}^3(p, \vec{w}) \rangle^{(0)} \\ &\quad \times \langle \epsilon \psi_{\vec{\beta} \vec{h}}^3(p+1, \vec{w}) \epsilon \bar{\psi}_{\vec{\alpha}_2 \vec{f}_2}^3(y) \rangle^{(0)}, \end{aligned} \quad (\text{B.2})$$

where the structure of Γ^{+e^0} restricts the spin indices in $\vec{\beta}$ to be lower. Hence, the *product* structure of Eq. (B.2) is obtained if we take $\vec{\alpha}_1$ and $\vec{\alpha}_2$ with only lower indices. This formula is also *closed* meaning that the correlation on the l.h.s. of Eq. (B.2) is the same as the two correlations appearing on the r.h.s. It is important to stress that, if we start with a gauge invariant average with generic fields with an odd number of fermions,

$\langle H(x) \bar{L}(y) \rangle$, we do not need to make an a priori guess of what the baryon fields are. They emerge as fields to achieve the product structure, and is consistent with confinement as the baryon fields are gauge invariant local fields with three quarks.

As explained above, the product property in Eq. (B.2) leads to the existence of particles with isolated dispersion curves, after eliminating the redundancies due to linear dependencies.

As each αf takes four values, there are 64 possible $\vec{\alpha} \vec{f}$'s but, by identifying states which are related by the totally symmetric property, this number is reduced to 20. The list of 12 representatives out of the 20 states with their multiplicities in the intermediate sums of Eq. (B.2) is the following: $\vec{f} = (+, +, +), (-, -, -)$, $\vec{\alpha} = (+, +, +), (-, -, -)$, multiplicity 1; $\vec{f} = ((+, +, +), (-, -, -))$, $\vec{\alpha} = (+, +, -), (-, -, +)$, multiplicity 3; $\vec{f} = (+, +, -)$, $\vec{\alpha} = (+, +, +), (-, -, -)$, multiplicity 3; $\vec{f} = (-, -, +)$, $\vec{\alpha} = (+, +, +), (-, -, -)$, multiplicity 3. The other 8 states are given below.

Eight of the above 12 states with $\vec{f} = (+, +, +), (-, -, -)$ have $I = 3/2$ and $I_z = 3/2, -3/2$, respectively, and $s = \pm 3/2, \pm 1/2$. For the remaining four states, we use the fact (from properties of the C-G coefficients) that $I = 3/2, I_z = \pm 1/2$ states formed from tensor products of three isospin 1/2 states have equal weights and are symmetric under interchange of any of the individual isospins. In this way, these states with $\vec{\alpha} = (+, +, +), (-, -, -)$ have $I = 3/2, I_z = \pm 1/2$. Thus, we are left with 8 states, with $|I_z| = 1/2$ and $|s| = 1/2$, given by (including the respective multiplicities on the right)

- 1) $\vec{f} = (+, -, +)$, $\vec{\alpha} = (+, -, +)$, $I_z = \frac{1}{2}$, $s = \frac{1}{2}$, 3;
- 2) $\vec{f} = (-, +, +)$, $\vec{\alpha} = (+, -, +)$, $I_z = \frac{1}{2}$, $s = \frac{1}{2}$, 6;
- 3) $\vec{f} = (+, -, +)$, $\vec{\alpha} = (+, -, -)$, $I_z = \frac{1}{2}$, $s = -\frac{1}{2}$, 6;
- 4) $\vec{f} = (-, +, +)$, $\vec{\alpha} = (+, -, -)$, $I_z = \frac{1}{2}$, $s = -\frac{1}{2}$, 3;
- 5) $\vec{f} = (+, -, -)$, $\vec{\alpha} = (+, -, +)$, $I_z = -\frac{1}{2}$, $s = \frac{1}{2}$, 6;
- 6) $\vec{f} = (-, +, -)$, $\vec{\alpha} = (+, -, +)$, $I_z = -\frac{1}{2}$, $s = \frac{1}{2}$, 3;
- 7) $\vec{f} = (-, +, -)$, $\vec{\alpha} = (+, +, -)$, $I_z = -\frac{1}{2}$, $s = -\frac{1}{2}$, 3;
- 8) $\vec{f} = (+, -, -)$, $\vec{\alpha} = (-, +, -)$, $I_z = -\frac{1}{2}$, $s = -\frac{1}{2}$, 6.

We identify linear combinations of these 8 states which we show to have $I = 3/2$ and $I = 1/2$. The proton (neutron) is identified with the $I = 1/2, I_z = 1/2$ ($-1/2$) state.

Coupling the first two isospins of 1) and 2) (by antisymmetric linear combination), to give a zero isospin, than coupling the third isospin gives the p_+ state. Similarly, antisymmetric linear combination of 3) and 4), gives p_- . n_+ (n_-) is obtained by antisymmetric linear combination of 5) and 6) (7) and 8)). For the $I = 3/2, I_z = 1/2, s = 1/2$ state with $\vec{\alpha} = (+, -, +)$ take the

linear combination $1) + 2) + 2)$ and note that the second 2) is not changed by $\alpha_1 f_1 \leftrightarrow f_3 \alpha_3$. Thus, the resulting state is totally symmetric in any $\alpha_j f_j \leftrightarrow f_k \alpha_k$; hence it is a $I = 3/2$ state. Similarly, the remaining three states are obtained with $(I, I_z, s) = (3/2, 1/2, -1/2)$, $(3/2, -1/2, 1/2)$, $(3/2, -1/2, -1/2)$, and are identified with the linear combinations $3) + 3) + 4)$, $5) + 5) + 6)$ and $7) + 8) + 8)$, respectively.

The above 20 states are unnormalized ones. Using the twenty normalized states of Section 3.1, and their multiplicities, and rewriting the intermediate states of Eq. (B.2) in terms of normalized states, we obtain, for $x^0 < y^0$ and $\vec{w} \in \mathbb{Z}^3$,

$$\langle B_{II_z s}(x) \bar{B}_{II_z s'}(y) \rangle^{(3)} = - \sum_{\vec{w}, r} \langle B_{II_z s}(x) \bar{B}_{II_z r}(p, \vec{w}) \rangle^{(0)} \times \langle B_{II_z r}(p+1, \vec{w}) \bar{B}_{II_z s'}(y) \rangle^{(0)}. \quad (\text{B.3})$$

Similar considerations apply for $x^0 > y^0$, and we finally obtain the product structure for the two-baryon correlation of Eq. (3.10)

$$G_{\ell_1 \ell_2}^{(3)}(x, y) = - \sum_{\vec{w}, \ell} G_{\ell_1 \ell}^{(0)}(x, (p, \vec{w})) G_{\ell \ell_2}^{(0)}((p+1, \vec{w}), y). \quad (\text{B.4})$$

Furthermore, for $\kappa = 0$, using Wick's theorem in the $\tilde{\psi}$ fields, we obtain the normalization property, for any x ,

$$\langle B_{II_z s}(x) \bar{B}_{II_z s}(x) \rangle^{(0)} = -1.$$

To continue, we now use symmetries at the level of correlation functions to give a more complete treatment of the baryon dispersion curves than that appearing in Theorem 4 of Ref. [5]. Besides the ordinary time reversal $\mathcal{T}$, charge conjugation $\mathcal{C}$ and parity $\mathcal{P}$, which can be implemented as unitary (anti-unitary for time reversal) operators acting on the physical Hilbert space $\mathcal{H}$, we also have a time *reflection* symmetry $\mathcal{T}$ as well as the combination $\mathcal{CP}\mathcal{T}$.

We summarize these symmetries as follows:

- Time Reflection $\mathcal{T}$: The $\tilde{\psi}$ fields transform as $\psi_\alpha(x) \rightarrow A_{\alpha\beta} \psi_\beta(-x^0, \vec{x})$, $\bar{\psi}_\alpha(x) \rightarrow \bar{\psi}_\beta(-x^0, \vec{x}) B_{\beta\alpha}$ where $A = B = A^{-1} = \begin{pmatrix} 0 & -iI_2 \\ iI_2 & 0 \end{pmatrix}$, $f(g_{xy}) \rightarrow f(g_{\bar{x}\bar{y}})$, with $\bar{z} = (-z^0, \vec{z})$;
- Charge Conjugation $\mathcal{C}$: $\psi_\alpha(x) \rightarrow \bar{\psi}_\beta(x) A_{\beta\alpha}$, $\bar{\psi}_\alpha(x) \rightarrow B_{\alpha\beta} \psi_\beta(x)$, $A = -B = B^{-1} = \begin{pmatrix} 0 & i\sigma^2 \\ i\sigma^2 & 0 \end{pmatrix}$, $f(g_{xy}) \rightarrow f(g_{xy}^*)$;
- Parity $\mathcal{P}$: $\psi_\alpha(x) \rightarrow A_{\alpha\beta} \psi_\beta(x^0, -\vec{x})$, $\bar{\psi}_\alpha(x) \rightarrow \bar{\psi}_\beta(x^0, -\vec{x}) B_{\beta\alpha}$ where $A = B = A^{-1} = \gamma^0$, $f(g_{xy}) \rightarrow f(g_{\bar{x}\bar{y}})$, with $\bar{z} = (z^0, -\vec{z})$;
- Rotation r_3 of $\pi/2$ about e^3 : $\psi_\alpha(x) \rightarrow A_{\alpha\beta} \psi_\beta(x^0, x^2, -x^1, x^3)$, $\bar{\psi}_\alpha(x) \rightarrow \bar{\psi}_\beta(x^0, x^2, -x^1, x^3) B_{\beta\alpha}$ where $A = B^{-1} =$

$\text{diag}(e^{-i\pi/4}, e^{i\pi/4}, e^{-i\pi/4}, e^{i\pi/4})$, $f(g_{xy}) \rightarrow f(g_{\bar{x}\bar{y}})^*$, with $\bar{z} = (z^0, z^2, -z^1, z^3 \bar{z})$;

- Time Reversal $\mathcal{T}$: $\psi_\alpha(x) \rightarrow \bar{\psi}_\beta(-x^0, \vec{x}) A_{\beta\alpha}$, $\bar{\psi}_\alpha(x) \rightarrow B_{\alpha\beta} \psi_\beta(-x^0, \vec{x})$, $A = B = B^{-1} = \gamma^0$, $f(g_{xy}) \rightarrow [f(g_{xy})]^*$.

The above symmetries are taken to be order preserving, except for $\mathcal{C}$ and $\mathcal{T}$ which are order reversing. For all of them, except time reversal, the field average equals the transformed field average; for time reversal the transformed field average is the complex conjugate of the field average.

Applying the $\mathcal{PC}\mathcal{T}$ symmetry, we find the following relations, after making a half-integer shift in the temporal coordinate of the lattice,

$$\langle p_\pm(u) \bar{p}_\mp(v) \rangle = 0 \quad , \quad \langle p_+(u) \bar{p}_+(v) \rangle = \langle p_-(u) \bar{p}_-(v) \rangle, \quad (\text{B.5})$$

and the same for the n 's. As shown in Section 3, we also have $\langle p_\pm(u) \bar{n}_\mp(v) \rangle = 0 = \langle n_\pm(u) \bar{p}_\mp(v) \rangle$, etc. Setting $\bar{\phi}_s(u) \equiv \bar{B}_{\frac{3}{2}\frac{3}{2}s}(u)$, we also have $\langle \phi_{\frac{3}{2}}(u) \bar{\phi}_{-\frac{3}{2}}(v) \rangle = \langle \phi_{\frac{1}{2}}(u) \bar{\phi}_{-\frac{1}{2}}(v) \rangle = 0$, $\langle \phi_{\mp\frac{3}{2}}(u) \bar{\phi}_{\mp\frac{1}{2}}(v) \rangle = -\langle \phi_{\pm\frac{1}{2}}(u) \bar{\phi}_{\pm\frac{3}{2}}(v) \rangle$, $\langle \phi_{-\frac{3}{2}}(u) \bar{\phi}_{\frac{1}{2}}(v) \rangle = \langle \phi_{-\frac{1}{2}}(u) \bar{\phi}_{\frac{3}{2}}(v) \rangle$, $\langle \phi_{\frac{3}{2}}(u) \bar{\phi}_{\frac{3}{2}}(v) \rangle = \langle \phi_{-\frac{3}{2}}(u) \bar{\phi}_{-\frac{3}{2}}(v) \rangle$, and $\langle \phi_{\frac{1}{2}}(u) \bar{\phi}_{\frac{1}{2}}(v) \rangle = \langle \phi_{-\frac{1}{2}}(u) \bar{\phi}_{-\frac{1}{2}}(v) \rangle$ and, using r_3 , for $x_r \equiv (x^0, x^2, -x^1, x^3)$,

$$\begin{aligned} \langle \phi_{\frac{3}{2}}(u) \bar{\phi}_{\frac{1}{2}}(v) \rangle &= -i \langle \phi_{\frac{3}{2}}(u_r) \bar{\phi}_{\frac{1}{2}}(v) \rangle, \\ \langle \phi_{\frac{3}{2}}(u) \bar{\phi}_{-\frac{1}{2}}(v) \rangle &= -\langle \phi_{\frac{3}{2}}(u_r) \bar{\phi}_{-\frac{1}{2}}(v) \rangle, \\ \langle \phi_{\frac{3}{2}}(u) \bar{\phi}_{-\frac{3}{2}}(v) \rangle &= i \langle \phi_{\frac{3}{2}}(u_r) \bar{\phi}_{-\frac{3}{2}}(v) \rangle, \\ \langle \phi_{\frac{1}{2}}(u) \bar{\phi}_{-\frac{1}{2}}(v) \rangle &= -i \langle \phi_{\frac{1}{2}}(u_r) \bar{\phi}_{-\frac{1}{2}}(v) \rangle, \\ \langle \phi_{\frac{1}{2}}(u) \bar{\phi}_{-\frac{3}{2}}(v) \rangle &= -\langle \phi_{\frac{1}{2}}(u_r) \bar{\phi}_{-\frac{3}{2}}(v) \rangle, \\ \langle \phi_s(u) \bar{\phi}_s(v) \rangle &= \langle \phi_s(u_r) \bar{\phi}_s(v) \rangle. \end{aligned} \quad (\text{B.6})$$

The relations after Eq. (B.5) carry over to the two-baryon correlation G .

Using Eq. (B.5), the proton and neutron two-point functions are diagonal in spin, at all orders in κ . For spin 3/2, label the basis by 3/2, -3/2, 1/2, and -1/2, or, simply, 1, 2, 3, and 4, respectively. With this labeling, and remarking that time reversal $\mathcal{T}$ parity $\mathcal{P}$ and the spectral representation ensures self-adjointness, the two-point function matrix has the structure

$$\begin{pmatrix} a & 0 & c & d \\ 0 & a & \bar{d} & -\bar{c} \\ \bar{c} & d & b & 0 \\ \bar{d} & -c & 0 & b \end{pmatrix},$$

with multiplicity two eigenvalues

$$\mu_\pm = \{(b+a) \pm [(b-a)^2 + 4(|c|^2 + |d|^2)]^{1/2}\}/2.$$

The corresponding eigenvectors are $(c, \bar{d}, \mu_\pm - a, 0)$ and $(\mu_\pm - b, 0, \bar{c}, \bar{d})$. Note that these two linearly independent vectors have a singular limit when $c, d \rightarrow 0$, where

they become identical. [This limit corresponds to taking $\tilde{\Gamma}_{13}, \tilde{\Gamma}_{14} \rightarrow 0$ in Eq. (B.8) below.]

The same matrix structure is satisfied by its inverse matrix and, consequently, it also holds for the Fourier transforms $\tilde{G}_{ss'}(p^0 = i\chi, \vec{p})$ and $\tilde{\Gamma}_{ss'}(p^0 = i\chi, \vec{p})$, $\chi \in \mathbb{R}$. Using the r_3 symmetry given above, shows that these matrices are diagonal for $\vec{p} = \vec{0}$ as well as for $\vec{p} = (p^1 = 0, p^2 = 0, p^3)$, which is used in Ref. [5] to determine the one-baryon mass spectrum and to show that no mass splitting occurs up to and including order κ^6 . For $\vec{p} \neq \vec{0}$, recall that the one-baryon dispersion curves $w(\vec{p})$ are the solutions of $\det \tilde{\Gamma}(p^0 = iw(\vec{p}), \vec{p}) = 0$. From the above structure, omitting the $\vec{p}$ dependence, the determinant factorizes as

$$\det \tilde{\Gamma} = [\lambda_+ \lambda_-]^2, \quad (\text{B.7})$$

which gives two by two identical dispersion curves associated to the zeroes of

$$\lambda_{\pm} \equiv \frac{1}{2}[\tilde{\Gamma}_{11} + \tilde{\Gamma}_{33}] \pm \sqrt{\frac{1}{4}[\tilde{\Gamma}_{11} - \tilde{\Gamma}_{33}]^2 + |\tilde{\Gamma}_{13}|^2 + |\tilde{\Gamma}_{14}|^2}. \quad (\text{B.8})$$

For $\vec{p} = \vec{0}$, $\lambda_+ = \tilde{\Gamma}_{11}$ and $\lambda_- = \tilde{\Gamma}_{33}$, and if they are not equal, mass splitting occurs.

We now show how to find the solution for $\lambda_+ = 0$, the treatment for other solution being similar. Suppressing the κ dependence and putting $c_3(\vec{p}) = c_3 \sum_{j=1}^3 2 \cos p^j$, for $c_3 = -1/8$ and $c_{30} = -1$ (see Ref. [5]), we can write

$$\begin{aligned} \tilde{\Gamma}_{s_1 s_2}(\vec{p}) = & [-1 - c_3(\vec{p})\kappa^3 - c_{30}\kappa^3(e^{ip^0} + e^{-ip^0})] \\ & \times \delta_{s_1 s_2} + \sum'_{n \geq 0, \vec{x}} \Gamma_{s_1 s_2}(n, \vec{x}) e^{-i\vec{p} \cdot \vec{x}} \\ & \times [\delta_{n0} + (1 - \delta_{n0})(e^{ip^0 n} + e^{-ip^0 n})], \end{aligned}$$

where $\sum'$ means all terms of order κ^6 or higher are maintained, and we have used $\Gamma(x^0, \vec{x}) = \Gamma(-x^0, \vec{x})$ which follows from $G(x^0, \vec{x}) = G(-x^0, \vec{x})$ by the spectral representation of Eq. (3.11).

To bring the solution in p^0 to the origin as $\kappa \searrow 0$, as in Ref. [5], we make a nonlinear transformation by introducing the auxiliary function $H(w, \kappa) \equiv H(w, \kappa, \vec{p})$ and a variable w such that $H(w = -1 - c_3(\vec{p})\kappa^3 - c_{30}\kappa^3 e^{-ip^0}, \kappa) = \lambda_+$. $H(w, \kappa)$ is then defined by

$$\begin{aligned} H(w, \kappa) = & w + \frac{c_{30}\kappa^6}{1 + w + c_3(\vec{p})\kappa^3} + \frac{1}{2} \sum''_{n \geq 0, \vec{x}} \\ & \times (\Gamma_{11} + \Gamma_{33})(n, \vec{x}) e^{-i\vec{p} \cdot \vec{x}} \{ \delta_{n0} + (1 - \delta_{n0}) \\ & \times (-1)^n [(\frac{1+w+c_3(\vec{p})\kappa^3}{c_{30}\kappa^3})^n + (\frac{c_{30}\kappa^3}{1+w+c_3(\vec{p})\kappa^3})^n] \} \\ & + \sqrt{\zeta_1^2 + |\zeta_2|^2 + |\zeta_3|^2}, \end{aligned} \quad (\text{B.9})$$

where

$$\begin{aligned} \zeta_1 = & \frac{1}{2} \sum''_{n \geq 0, \vec{x}} (\Gamma_{33} - \Gamma_{11})(n, \vec{x}) e^{-i\vec{p} \cdot \vec{x}} \{ \delta_{n0} + (1 - \delta_{n0}) \\ & \times (-1)^n [(\frac{1+w+c_3(\vec{p})\kappa^3}{c_{30}\kappa^3})^n + (\frac{c_{30}\kappa^3}{1+w+c_3(\vec{p})\kappa^3})^n] \}, \end{aligned}$$

and similarly for ζ_2 and ζ_3 with the factor $[\Gamma_{33} - \Gamma_{11}]/2$ replaced by Γ_{13} and Γ_{14} , respectively. In the above, $\sum''$

means that we only retain terms of order κ^6 or higher in the sum, and that we have used the short distance behavior of $\Gamma_{s_1 s_2}$ as given in Theorem 2 of Ref. [5] [the misprinted values of c_3 and c_{30} in this reference are corrected above].

Before we continue the analysis, we give the following short distance behavior that extends the results of Theorem 2 of Ref. [5] to a higher order in κ (c_6 here is a κ -independent computable constants):

$$G_{s_1 s_2}(x) = \begin{cases} s_1 = s_2 = \frac{3}{2} \begin{cases} -4c_6\kappa^6, x = (0, 1, 1, 0); \\ 2c_6\kappa^6, x = (0, 1, 0, 1), \\ (0, 0, 1, 1); \end{cases} \\ s_1 = s_2 = \frac{1}{2} \begin{cases} 4c_6\kappa^6, x = (0, 1, 1, 0); \\ -2c_6\kappa^6, x = (0, 1, 0, 1), \\ (0, 0, 1, 1); \end{cases} \\ 0 \kappa^6, s_1 \neq s_2. \end{cases} \quad (\text{B.10})$$

We also find the following κ^9 contributions ($c_9, c_{9,\pm}$ being nonzero constants):

$$G_{s_1 s_2}(x) = \begin{cases} c_9\kappa^9, x = (0, 1, 1, 1), s_1 = s_2, \\ c_{9,\pm}\kappa^9, x = (0, 1, 1, 1), s_1 = \frac{3}{2}, s_2 = \pm\frac{1}{2}. \end{cases} \quad (\text{B.11})$$

Note that there are $\mathcal{O}(\kappa^8)$ contributions to $\mathcal{G}_{s_1 s_2}(x = 0)$ arising from two square loops starting and ending at $x = 0$, but with opposite orientations. They give equal diagonal contributions and there are *no* off-diagonal contributions using the symmetry properties of Eq. (B.6).

If any two of the terms in the square root are zero, the square root in Eq. (B.9) disappears and, since by Eq. (B.9) and the decay properties of Γ obtained with the hyperplane method leads to a joint analyticity in κ, w and each $\vec{p}$ component, we can apply the analytic implicit function theorem to find an analytic $w(\vec{p}, \kappa)$ such that $H(w(\kappa, \vec{p}), \kappa, \vec{p}) = 0$, for small κ . In fact, the analytic implicit function $w(\vec{p}, \kappa)$ becomes explicit and admits the contour integral representation

$$w(\vec{p}, \kappa) = \frac{1}{2\pi i} \int_{|w|=r} \frac{w(\partial H / \partial w)(w, \kappa, \vec{p})}{H(w, \kappa, \vec{p})} dw,$$

where the contour can be taken as a small positively oriented circle (see, for instance, Ref. [47]). This representation is used to obtain an explicit formula for the Taylor series coefficients in Ref. [48].

Concerning the proton and the neutron, we know that their dispersion curves are identical by isospin symmetry. Moreover, as the $\mathcal{PC}$ T symmetry [see Eq. (B.5)] ensures $\tilde{\Gamma}_{s_1 s_2}(\vec{p})$ is diagonal (even at $\vec{p} \neq \vec{0}$) and proportional of the identity, the proton dispersion curves are also identical for their two spin states. The resulting proton and neutron dispersion curves are given by, recalling that $p_\ell^2 \equiv 2 \sum_{i=1}^3 (1 - \cos p^i)$ as in Eq. (3.14),

$$w(\vec{p}) \equiv w(\vec{p}, \kappa) = [-3 \ln \kappa - 3\kappa^3/4 + p_\ell^2 \kappa^3/8] + r_j(\vec{p}), \quad (\text{B.12})$$

where $r(\vec{p})$ is of $\mathcal{O}(\kappa^6)$ and is jointly analytic for each $\vec{p}$ component, and κ . As easily seen, $w(\vec{p})$ in Eq. (B.12)

is convex for small $|\vec{p}|$ and $|\kappa|$, and increases with $|\vec{p}|$, as physically expected.

As seen in Eq. (B.11), we have non-diagonal contributions for $G_{s_1 s_2}$, $|s_1| \neq |s_2|$. The square root in Eq. (B.9) survives. Using r_3 symmetry through Eq. (B.6) and $\mathcal{P}$, we find the following $\mathcal{O}(\geq \kappa^6)$ contributions to $\tilde{G}_{s_1 s_2}(\vec{p})$:

$$\begin{aligned}\tilde{G}_{\frac{3}{2}\frac{3}{2}}(\vec{p}) &\propto (-2 \cos p^1 \cos p^2 + \cos p^1 \cos p^3, \\ &\quad + \cos p^2 \cos p^3) \kappa^6, \\ \tilde{G}_{\frac{1}{2}\frac{1}{2}}(\vec{p}) &\propto (2 \cos p^1 \cos p^2 - \cos p^1 \cos p^3, \\ &\quad - \cos p^2 \cos p^3) \kappa^6, \quad (\text{B.13}) \\ \tilde{G}_{\frac{3}{2}\frac{1}{2}}(\vec{p}) &\propto [(1-i) \sin p^1 \cos p^2 \\ &\quad - (1+i) \cos p^1 \sin p^2] \sin p^3 \kappa^9, \\ \tilde{G}_{\frac{1}{2}\frac{1}{2}}(\vec{p}) &\propto \sin p^1 \sin p^2 \cos p^3 \kappa^9,\end{aligned}$$

where we observe that *all* of these contributions do vanish at $\vec{p} = \vec{0}$.

We have similar expressions for $\tilde{\Gamma}_{ij}$. Hence, $\tilde{\Gamma}_{\frac{3}{2}\frac{3}{2}}(\vec{p}) - \tilde{\Gamma}_{\frac{1}{2}\frac{1}{2}}(\vec{p}) \neq 0$, and we cannot successfully apply the same method as before, as the κ analyticity region shrinks to zero as $\vec{p} \searrow \vec{0}$. Further analysis is then required.

Using the above results, without using analyticity, we can however obtain the remaining approximate dispersion curves, but with less knowledge about their behavior. Particularly, we only have a rough bound on $r(\vec{p})$. To do this job, with h_2 denoting the square root term in Eq. (B.9), we can write

$$H = w + h_1 + h_2, \quad (\text{B.14})$$

where, uniformly in $\vec{p}$, we have $|h_1| \leq c_1 \kappa^6$ and $|h_2| \leq c_2 \kappa^6$, for some positive constants $c_{1,2}$, by the above and the bounds of Theorem 2 of Ref. [5]. [The h_2 bound is due to the κ^6 term in ζ_1 , which goes to zero as $\vec{p} \searrow \vec{0}$, but we can possibly have a term of order $\mathcal{O}(\geq \kappa^{10})$ which does not go to zero as $\vec{p} \searrow \vec{0}$. Such a term leads to a mass splitting.]

Writing $H = 0$ in Eq. (B.14) as

$$w = -h_1(w, \kappa, \vec{p}) - h_2(w, \kappa, \vec{p}), \quad |h_1 + h_2| \leq c \kappa^6,$$

we plot the left and right sides as functions of w . For $|w| \leq (1 + \alpha) c \kappa^6$, for some $\alpha > 0$, we have at least one solution of $H = 0$. Similar considerations apply in working out the zero determinant solution for λ_- in Eq. (B.7). From Ref. [5], a Rouché theorem argument shows that there are exactly four solutions of $\det \tilde{\Gamma} = 0$, so that there is exactly one solution for each $\lambda_{\pm}$, which are presented in Eq. (3.15).

We now give an explicit construction of the dispersion curve in the non-diagonal case. In $H(w, \kappa, \vec{p})$, it is convenient to make another nonlinear variable transformation from w to $u \equiv w + f_1(w, \kappa)$ and, by the analytic implicit function theorem, $L(u, w) \equiv w + w + f_1(w, \kappa) - u = 0$ yields the analytic function $w \equiv r(u, \kappa)$ such that

$L(r(u, \kappa), u) = 0$ for $|w|, |\kappa|$ small. The condition for the existence of r is satisfied as

$$(\partial L / \partial w)(w, u) = 1 + (\partial f_1 / \partial w)(w, \kappa) \neq 0,$$

for $|w|, |u|, |\kappa|$ small, since $|f_1| \leq c_1 \kappa^6$ and $|\partial f_1 / \partial w| \leq c_2 \kappa^6$. Furthermore, $(\zeta_1^2 + |\zeta_2|^2 + |\zeta_3|^2)(w = r(u, \kappa), \kappa, \vec{p})$ is jointly analytic in u, κ, p^j and we write it as $\sum_n a_n u^n$.

Thus, we can write the dispersion curve determining equation $H = 0$ as the fixed point equation

$$u = \left[\sum_n a_n u^n \right]^{1/2} \equiv f(u).$$

We obtain the solution $\underline{u}$ by iteration. The $(\ell - 1)$ th iteration gives

$$\begin{aligned}u &= \left[\sum_{n_1} a_{n_1} \left[\sum_{n_2} a_{n_2} \left[\sum_{n_3} a_{n_3} \dots \left[\sum_{n_\ell} a_{n_\ell} u^{n_\ell} \right]^{\frac{(n_\ell - 1)}{2}} \right. \right. \right. \\ &\quad \times \dots \left. \left. \left. \right]^{\frac{n_2}{2}} \right]^{\frac{n_1}{2}} \right]^{1/2}.\end{aligned}$$

Assuming convergence to the fixed point $\underline{u}$, for the initial value $u = 0$, we take the limit $\ell \rightarrow \infty$ in the above, with a_ℓ replaced by $a_0 \delta_{\ell 0}$, to get the solution $\underline{u}$.

Additionally, by adapting the subtraction method of Ref. [5], it can be shown that the isolated dispersion curves (the upper gap property) associated with the two-baryon functions of the above 20 particles, with asymptotic mass $-3 \ln \kappa$, are the only particles in the full space of states generated by an odd number of ψ fields, up to the meson-baryon threshold $\approx -5 \ln \kappa$ (the meson asymptotic masses are $\approx -2 \ln \kappa$).

To close this Appendix, we point out that for free lattice Fermi field the matrix structure of the Fourier transform of the two-point function is the same as in the continuum, i.e. $p^\mu \gamma^\mu + m I_2$ in the lower subspace.

From this explicit relation, we see that there is *no* spin flip and that the diagonal elements are the same. Another way to see this is to use the $\mathcal{PCT}$ symmetry which gives $\langle \psi_3 \psi_4 \rangle = \langle \psi_4 \psi_3 \rangle = 0$ and $\langle \psi_3 \psi_3 \rangle = \langle \psi_4 \psi_4 \rangle$. Moreover, in the interacting case with a rotationally invariant interaction, such as in the $U(N)$ -invariant Gross-Neveu model, the same holds using $\mathcal{PCT}$ symmetry.

However, for the three-quark composite local baryon fields, the $\mathcal{PCT}$ symmetry, or even the $\mathcal{PC}$ T symmetry, are not enough to conclude diagonality of the two-point function in the spin, except for protons and neutrons. As we have seen, it is not diagonal in the spin 3/2 case.

In the continuum case, relativistic invariance allows us to pass to the rest frame, and the two-point function transforms according to the spatial rotation group $SU(2)$. There is no (spin) mass splitting and there is just one dispersion curve, the relativistic one.

APPENDIX C: Coincident point four-baryon correlation and linear dependence relation

Here we obtain the decomposition of $M_0^{(0)}$ in the total

spin basis, and show the pointwise linear dependence relation given in Section 4.3 [see Eq. (4.27)].

At $\kappa = 0$, using the total spin spatial symmetry and orthogonality properties, the following block decomposition as given in Eq. (4.25) is

$$M^{(0)}(0) = M_1^{(0)}(0) \oplus M_1^{(0)}(0) \oplus M_1^{(0)}(0) \oplus M_b^{(0)}(0).$$

It is convenient to use the shorthand notation $b_{I_z s} \equiv B_{(I)I_z s}$. $M_b^{(0)}(0)$ is diagonal by the total spin orthogonality relation and, by the lowering relations of the z -component of total spin, all of its elements are equal. The first diagonal element is

$$\begin{aligned} \langle T_{T=3T_z=3}^{3/2\,3/2} \bar{T}_{T=3T_z=3}^{3/2\,3/2} \rangle^{(0)} &= 4 \langle b_{3/2\,3/2} b_{-3/2\,3/2} \\ &\quad \times \bar{b}_{3/2\,3/2} \bar{b}_{-3/2\,3/2} \rangle^{(0)} = -4. \end{aligned}$$

The elements of the 2×2 matrix $M_1^{(0)}(0)$, with $S = 1$, $S_z = 1$, $T = 1$, $T_z = 1$, are given by

$$\begin{aligned} \mu_{11} &= 2 \langle p_{+n} + \bar{p}_{+n} \rangle^{(0)}, \quad \mu_{22} = \langle \chi \bar{\chi} \rangle^{(0)}, \\ \mu_{21} &= \mu_{12} = \langle p_{+n} + \bar{\chi} \bar{\chi} \rangle^{(0)}, \end{aligned}$$

where $\chi \equiv \sqrt{3/10} b_{3/2\,3/2} b_{-3/2\,-1/2} - \sqrt{2/5} b_{3/2\,1/2} b_{-3/2\,1/2} + \sqrt{3/10} b_{3/2\,-1/2} b_{-3/2\,3/2} - \sqrt{3/10} b_{1/2\,3/2} b_{-1/2\,-1/2} + \sqrt{2/5} b_{1/2\,1/2} b_{-1/2\,1/2} - \sqrt{3/10} b_{1/2\,-1/2} b_{-1/2\,3/2}$.

For the sake of checking the calculations, we compute the term of the four-baryon correlation appearing in μ_{ij} with b fields which have the prefactor 1 since their values are integers. We denote them by B^u and call them unnormalized fields, and label the unnormalized products p_{+n}^u , $b_{\frac{3}{2}\,\frac{3}{2}}^u b_{\frac{-3}{2}\,\frac{-1}{2}}^u$, $b_{\frac{3}{2}\,\frac{1}{2}}^u b_{\frac{-3}{2}\,\frac{1}{2}}^u$, $b_{\frac{3}{2}\,\frac{-1}{2}}^u b_{\frac{-3}{2}\,\frac{3}{2}}^u$, $b_{\frac{1}{2}\,\frac{3}{2}}^u b_{\frac{-1}{2}\,\frac{-1}{2}}^u$, $b_{\frac{1}{2}\,\frac{1}{2}}^u b_{\frac{-1}{2}\,\frac{1}{2}}^u$, $b_{\frac{1}{2}\,\frac{-1}{2}}^u b_{\frac{-1}{2}\,\frac{3}{2}}^u$ appearing above by 1, 2, ..., 7, respectively. Here, the corresponding $\kappa = 0$ averages are denoted by (i, j) , where the second spot corresponds to the product of two barred fields. By isospin flip symmetry, we have the following equalities: $(1, 2) = (1, 4)$, $(1, 5) = (1, 7)$, $(2, 2) = (4, 4)$, $(2, 6) = (4, 6)$, $(3, 5) = (3, 7)$, and $(5, 5) = (5, 7)$. Thus, all we need to calculate are the elements listed below with their respective values:

$$\begin{aligned} (1, 1) &= -360, \quad (1, 2) = 144, \quad (1, 3) = -96 \\ (1, 5) &= -144, \quad (2, 6) = 288, \quad (3, 3) = -144 \\ (3, 5) &= 96, \quad (3, 6) = 240, \quad (4, 7) = 144 \\ (5, 5) &= -240, \quad (5, 7) = -192, \quad (6, 6) = -1296. \end{aligned}$$

Reinstating the normalization prefactors and substituting in μ_{ij} results in the matrix $M_1^{(0)}(0)$ of Eq. (4.26).

We now turn to the linear dependence relation of Eq. (4.27). To prove that

$$\mathcal{T}_{S=1, S_z} + \frac{\sqrt{5}}{2} \mathcal{T}_{T=1, T_z} = 0,$$

for $S_z = T_z = -1, 0, 1$, it is enough to show it holds for $S_z = T_z = 1$ and apply the spin lowering operator

on S_z and T_z . For $S_z = T_z = 1$, using the definition of the $\mathcal{T}$ fields in Section 4.1 [see Eq. (4.20)], we write $\mathcal{T}_{S=1,1} + \frac{\sqrt{5}}{2} \mathcal{T}_{T=1,1}$ as the sum, with from now on $b_{I_z s} \equiv B_{(I=3/2)I_z s}$, $s = \pm 3/2, \pm 1/2$,

$$\begin{aligned} &\sqrt{2} p_{+n} + \frac{\sqrt{5}}{2} \left[\sqrt{\frac{3}{8}} b_{\frac{3}{2}\,\frac{3}{2}} b_{\frac{-3}{2}\,\frac{-1}{2}} - \sqrt{\frac{1}{2}} b_{\frac{3}{2}\,\frac{1}{2}} b_{\frac{-3}{2}\,\frac{1}{2}} \right. \\ &+ \sqrt{\frac{3}{8}} b_{\frac{3}{2}\,\frac{-1}{2}} b_{\frac{-3}{2}\,\frac{3}{2}} - \sqrt{\frac{3}{8}} b_{\frac{1}{2}\,\frac{3}{2}} b_{\frac{-1}{2}\,\frac{-1}{2}} + \sqrt{\frac{1}{2}} b_{\frac{1}{2}\,\frac{1}{2}} b_{\frac{-1}{2}\,\frac{1}{2}} \\ &\quad \left. - \sqrt{\frac{3}{8}} b_{\frac{1}{2}\,\frac{-1}{2}} b_{\frac{-1}{2}\,\frac{3}{2}} \right]. \end{aligned}$$

A typical term in the above is labeled by a product of six ψ 's, with indices from one of the following 12 sets of indices: (1) $1 - -$; (2) $1 - +$; (3) $1 + -$; (4) $1 + +$; (5) $2 - -$; (6) $2 - +$; (7) $2 + -$; (8) $2 + +$; (9) $3 - -$; (10) $3 - +$; (11) $3 + -$; (12) $3 + +$.

There can be no repeats of any of the twelve indices in the six factors, by Pauli exclusion. We are left with 21 linearly independent products of six ψ 's with no repeated indices, and the coefficients of each of these linearly independent elements sum to zero.

APPENDIX D: Decay of the Bethe-Salpeter kernel

Using the hyperplane decoupling method, as described in Appendix B (see also Ref. [5] for more details), we obtain improved temporal decay for the kernel K , as compared to that of M and M_0 . For now, we use the original and unmodified M , and we will treat the modified $M \equiv M'$ at the end of the Appendix.

The temporal decay follows from a product structure of the sixth derivatives of the four-point functions M and M_0 , which in turn follows from the product structure of the sixth κ_p derivative of the basic four-point functions D and D_0 defined in Section 4.2. Recall that D , in the individual spin basis is defined in Eq. (4.22) and D_0 is obtained from D by erroneously applying Wick's theorem to the baryon fields in D . In a similar way, M_0 is obtained from M . In what follows, we suppress the collective index subscripts from D and D_0 . The product structure extends from D (D_0) to M (M_0) as it is preserved under real orthogonal transformations.

We show that, for $x_1^0 = x_2^0 \leq p < x_3^0 = x_4^0$, and with $\vec{w}, \vec{w}_1, \vec{w}_2 \in \mathbb{Z}^3$,

$$\begin{aligned} D^{(6)}(x_1, x_2, x_3, x_4) &= -\frac{1}{2} \sum_{\vec{w}_1 \neq \vec{w}_2} \\ &\quad \times D^{(0)}(x_1, x_2, (p, \vec{w}_1), (p, \vec{w}_2)) \\ &\quad \times D^{(0)}((p+1, \vec{w}_1), (p+1, \vec{w}_2), x_3, x_4) \\ &\quad - \frac{1}{4} \sum_{\vec{w}} D^{(0)}(x_1, x_2, (p, \vec{w}), (p, \vec{w})) \\ &\quad \times D^{(0)}((p+1, \vec{w}), (p+1, \vec{w}), x_3, x_4), \end{aligned} \quad (D1)$$

and

$$\begin{aligned} D_0^{(6)}(x_1, x_2, x_3, x_4) &= -\frac{1}{2} \sum_{\vec{w}_1, \vec{w}_2} \\ &\quad \times D_0^{(0)}(x_1, x_2, (p, \vec{w}_1), (p, \vec{w}_2)) \\ &\quad \times D_0^{(0)}((p+1, \vec{w}_1), (p+1, \vec{w}_2), x_3, x_4), \end{aligned} \quad (D2)$$

and similarly for $x_1^0 > x_3^0$. We write these results schematically as

$$D^{(6)} = -\frac{1}{2}D^{(0)} \circledast D^{(0)} - \frac{1}{4}D^{(0)} \odot D^{(0)}, \quad (D3)$$

and

$$D_0^{(6)} = -\frac{1}{2}D^{(0)} \circ D^{(0)}, \quad (D4)$$

using the convolution-like operations “ $\circledast$ ”, “ $\odot$ ” and “ $\circ$ ” given by Eqs. (D1) and (D2).

We now show how this structure gives the improved temporal decay for $K = D_0^{-1} - D^{-1}$. Take $x_1^0 \leq p < x_3^0$, for example, and let Λ (Λ_0) denote the inverse of D (D_0). It is easily seen that $K^{(n)} = 0$, $n = 0, 1, \dots, 5$, which follows from the fact that the same holds for D and D_0 . This is seen by expanding the numerators and using imbalance of fermions. We have, using Eqs. (D3) and (D4),

$$\begin{aligned} K^{(6)} &= \Lambda_0^{(0)} D_0^{(6)} \Lambda_0^{(0)} - \Lambda^{(0)} D^{(6)} \Lambda^{(0)} \\ &= +\Lambda_0^{(0)} (-\frac{1}{2}D_0^{(0)} \circ D_0^{(0)}) \Lambda_0^{(0)} \\ &\quad - \Lambda^{(0)} (-\frac{1}{2}D^{(0)} \circledast D^{(0)}) \Lambda^{(0)} \\ &\quad - \Lambda^{(0)} (-\frac{1}{4}D^{(0)} \odot D^{(0)}) \Lambda^{(0)} = 0, \end{aligned} \quad (D5)$$

for $x_1^0 + 1 < x_3^0$. As this holds for each hyperplane between x_1^0 and x_3^0 , the above leads to a $\kappa^6 \kappa^{7(x_3^0 - x_1^0 - 1)}$ decay, for $x_3^0 > x_1^0 + 1$ and a decay κ^6 , for $x_3^0 = x_1^0 + 1$. The κ^7 can be further improved to κ^8 taking into account fermion imbalance (or gauge integration) in the calculation of the seventh κ_p derivative.

We return to show how Eqs. (D1) and (D2) are obtained. For $x_1^0 = x_2^0 < x_3^0 = x_4^0$,

$$\begin{aligned} D_0^{(6)} &= -\langle B_{\ell_1}(x_1) \bar{B}_{\ell_3}(x_3) \rangle^{(3)} \langle B_{\ell_2}(x_2) \bar{B}_{\ell_4}(x_4) \rangle^{(3)} \\ &\quad + \langle B_{\ell_1}(x_1) \bar{B}_{\ell_4}(x_4) \rangle^{(3)} \langle B_{\ell_2}(x_2) \bar{B}_{\ell_3}(x_3) \rangle^{(3)}, \end{aligned}$$

where the ℓ 's are collective indices.

Using Eq. (B.3), and regrouping the terms, Eq. (D2) follows.

To show Eq. (D1), in addition to the gauge integral $\mathcal{I}_3$ for three coincident bonds of Eq. (B.1), we need to compute the gauge integral $\mathcal{I}_6$ of six coincident bonds (same orientation!). To compute $\mathcal{I}_6$, we use the generating function given in Refs. [2, 5] and differentiate with respect to the external sources. It is given by, with $U(g) \equiv g$ and

$$U^{-1}(g) \equiv g^{-1},$$

$$\begin{aligned} \mathcal{I}_6 &= \int g_{a_1 b_1} g_{a_2 b_2} g_{a_3 b_3} g_{a_4 b_4}^{-1} g_{a_5 b_5}^{-1} g_{a_6 b_6}^{-1} d\mu(g) \\ &= \frac{2}{3!4!} [\epsilon_{a_1 a_2 a_3} \epsilon_{b_1 b_2 b_3} \epsilon_{a_4 a_5 a_6} \epsilon_{b_4 b_5 b_6} + \\ &\quad \epsilon_{a_1 a_2 a_4} \epsilon_{b_1 b_2 b_4} \epsilon_{a_3 a_5 a_6} \epsilon_{b_3 b_5 b_6} + \\ &\quad \epsilon_{a_1 a_2 a_5} \epsilon_{b_1 b_2 b_5} \epsilon_{a_3 a_4 a_6} \epsilon_{b_3 b_4 b_6} + \\ &\quad \epsilon_{a_1 a_2 a_6} \epsilon_{b_1 b_2 b_6} \epsilon_{a_3 a_4 a_5} \epsilon_{b_3 b_4 b_5} + \\ &\quad \epsilon_{a_1 a_3 a_4} \epsilon_{b_1 b_3 b_3} \epsilon_{a_2 a_5 a_6} \epsilon_{b_2 b_5 b_6} + \\ &\quad \epsilon_{a_1 a_3 a_5} \epsilon_{b_1 b_3 b_5} \epsilon_{a_2 a_4 a_6} \epsilon_{b_2 b_4 b_6} + \\ &\quad \epsilon_{a_1 a_3 a_6} \epsilon_{b_1 b_3 b_6} \epsilon_{a_2 a_4 a_5} \epsilon_{b_2 b_4 b_5} + \\ &\quad \epsilon_{a_1 a_4 a_5} \epsilon_{b_1 b_4 b_5} \epsilon_{a_2 a_3 a_6} \epsilon_{b_2 b_3 b_6} + \\ &\quad \epsilon_{a_1 a_4 a_6} \epsilon_{b_1 b_4 b_6} \epsilon_{a_2 a_3 a_5} \epsilon_{b_2 b_3 b_5} + \\ &\quad \epsilon_{a_1 a_5 a_6} \epsilon_{b_1 b_5 b_6} \epsilon_{a_2 a_3 a_4} \epsilon_{b_2 b_3 b_4}]. \end{aligned} \quad (D6)$$

By performing appropriate (index) contractions in $\mathcal{I}_6$, we verify that $\mathcal{I}_4$, $\mathcal{I}_3$ and $\mathcal{I}_2$ are obtained, where (see Refs. [2, 5, 10])

$$\mathcal{I}_2 = \int g_{a_1 b_1} g_{a_2 b_2}^{-1} d\mu(g) = \frac{1}{3} \delta_{a_1 b_2} \delta_{a_2 b_1}, \quad (D7)$$

$\mathcal{I}_3$ is given in Eq. (B.1) and

$$\begin{aligned} \mathcal{I}_4 &= \int g_{a_1 b_1} g_{a_2 b_2}^{-1} g_{a_3 b_3} g_{a_4 b_4}^{-1} d\mu(g) \\ &= \frac{1}{8} [\delta_{a_1 b_2} \delta_{a_3 b_4} \delta_{b_1 a_2} \delta_{b_3 a_4} + (a_2 \rightleftharpoons a_4, b_2 \rightleftharpoons b_4)] \\ &\quad - \frac{1}{24} [\delta_{a_1 b_2} \delta_{a_3 b_4} \delta_{b_1 a_4} \delta_{b_3 a_2} + (a_2 \rightleftharpoons a_4, b_2 \rightleftharpoons b_4)]. \end{aligned} \quad (D8)$$

Eq. (D6) can be written in a compact notation as

$$\mathcal{I}_6 = \frac{2}{3!4!} \sum_{\mathcal{P}} E_R E_S,$$

where $\mathcal{P}$ is the set of the ten disjoint partitions of $T \equiv \{1, 2, 3, 4, 5, 6\} = R \cup S$, $|R| = 3 = |S|$, and for $R = \{i, j, k\}$, $E_R = \epsilon_{a_i a_j a_k} \epsilon_{b_i b_j b_k}$, where the elements in E_R and E_S are written in increasing order.

$D^{(6)}$ is obtained by writing $D = N/Z$ and expanding the exponential in the numerator of N . In this way, $D^{(6)} = N^{(6)}/Z^{(0)}$, where $N^{(6)} = N_n^{(6)} + N_o^{(6)}$. $N_n^{(6)}$ arises from the terms where there are two nonoverlapping bonds, indexed by w_1 and w_2 , each bond with three positively oriented gauge fields. After performing the gauge integrals using the formula for $\mathcal{I}_3$ of Eq. (B.1), we get

$$\begin{aligned} N_n^{(6)} &= -\frac{1}{2 \times 6^4} \epsilon_{a_1 a_2 a_3} \epsilon_{a_4 a_5 a_6} \epsilon_{b_1 b_2 b_3} \epsilon_{b_4 b_5 b_6} \sum_{\vec{w}_1 \neq \vec{w}_2, \{\alpha_i\}, \{f_i\}} \\ &\quad \times \langle B_{\ell_1}(x_1) B_{\ell_2}(x_2) (\bar{\psi}_{a_1 \alpha_1 f_1} \bar{\psi}_{a_2 \alpha_2 f_2} \bar{\psi}_{a_3 \alpha_3 f_3})(p, \vec{w}_1) \\ &\quad \times (\bar{\psi}_{a_4 \alpha_4 f_4} \bar{\psi}_{a_5 \alpha_5 f_5} \bar{\psi}_{a_6 \alpha_6 f_6})(p, \vec{w}_2) \rangle^{(0)} \\ &\quad \times \langle (\psi_{b_1 \alpha_1 f_1} \psi_{b_2 \alpha_2 f_2} \psi_{b_3 \alpha_3 f_3})(p+1, \vec{w}_1) \\ &\quad \times \psi_{a_4 \alpha_4 f_4} \psi_{a_5 \alpha_5 f_5} \psi_{a_6 \alpha_6 f_6})(p+1, \vec{w}_2) \\ &\quad \times \bar{B}_{\ell_3}(x_3) \bar{B}_{\ell_4}(x_4) \rangle^{(0)}, \end{aligned}$$

where the numerical factor comes from $(-1/2)^6 \times (1/6!) \times 20 \times (-2)^6 \times (1/2) \times (1/6)^2$; 20 is a combinatorial factor, $(-2)^6$ comes from the six Γ^{e^0} 's and $(1/6)^2$ coming from the two integrals $\mathcal{I}_3$. We observe that the sums over the α_i 's are restricted to lower components $\alpha = 3, 4$ due to the structure of Γ^{e^0} .

Resumming over the intermediate states, after taking into account multiplicities and normalizations of the baryon fields, gives the first term of Eq. (D1).

The contribution to N_o^6 , for which there are overlapping bonds, after performing the $\mathcal{I}_6$ gauge integral, is

$$\begin{aligned} & -\frac{1}{4 \times 6^4 \times 10} \sum_{\vec{w}, \{\alpha_i\}, \{f_i\}} [\epsilon_{a_1 a_2 a_3} \epsilon_{b_1 b_2 b_3} \epsilon_{a_4 a_5 a_6} \epsilon_{b_4 b_5 b_6} \\ & + \text{nine remaining terms}] \langle B_{\ell_1}(x_1) B_{\ell_2}(x_2) \\ & \times (\bar{\psi}_{a_1 \alpha_1 f_1} \dots \bar{\psi}_{a_6 \alpha_6 f_6})(p, \vec{w}) \rangle^{(0)} \\ & \times \langle (\psi_{b_1 \alpha_1 f_1} \dots \psi_{b_6 \alpha_6 f_6})(p+1, \vec{w}) \bar{B}_{\ell_3}(x_3) \bar{B}_{\ell_4}(x_4) \rangle^{(0)}, \end{aligned}$$

where the numerical factor comes from $(-1/2)^6 \times (1/6!) \times (-2)^6 \times [2/(3! \times 4!)]$; the term in square brackets is the contribution of the $\mathcal{I}_6$ integral. Again, resumming over the intermediate states gives the second term of Eq. (D1). In both cases, a factor 6^{-4} is absorbed in the normalization process.

By considering hyperplanes in each spatial direction, and additional complex parameters associated with them, spatial decay of the B-S kernel can also be obtained as in Ref. [26].

We now describe how the product structure of Eqs. (D3) and (D4) extend to $M^{(6)}$ and $M_0^{(6)}$. First, we take isospin C-G linear combinations of the external fields present in $D^{(6)}$, expressed in the individual spin and isospin basis, to get total isospin zero states. Then, again using the inverse C-G coefficients, we express the intermediate states in terms of states of fixed total isospin I and its z -components I_z . At this point, the isospin orthogonality properties are used to get only zero total isospin intermediate states. In this way, we arrive at the following product structure formula for $M^{(6)}$

$$M^{(6)} = -\frac{1}{2} M^{(0)} \otimes M^{(0)} - \frac{1}{4} M^{(0)} \odot M^{(0)},$$

where we remember that we are still in the individual spin basis. We can pass to the total isospin zero, total spin basis by taking spin C-G linear combinations. The product structure is maintained as it is invariant under real orthogonal transformations. A similar argument applies to $M_0^{(6)}$.

We are now ready to explain and motivate the adoption of a h -modified $M \equiv M'$ as introduced in Section 4.3. For this, we consider the following leading contributions to the unmodified $K(x_1, x_2, x_3, x_4)$ and the corresponding contributions to the generalized potential $\hat{K}(\vec{\xi}, \vec{\eta}, k^0)$: *i*) the κ^0 order, $K^{(0)}$ at coincident points, and corresponding space-range zero, energy independent constant potential $\hat{K}(\vec{0}, \vec{0}, k^0)$; *ii*) the κ^6 contribution with time-range one and space-range zero $K^{(6)}(0, 0, e^0, e^0)$, with energy-dependent zero space-range potential $\hat{K}(0, 0, k^0 = i(2\bar{m} -$

$\epsilon))$ of order $\kappa^2 e^{-\epsilon}$; and, *iii*) the κ^2 quasimeson exchange κ^2 contribution of space-range one $K^{(2)}(0, 0, e^j, e^j)$, $j = 1, 2, 3$, with space-range one energy-independent potential $\hat{K}(0, e^j, k^0)$ of order κ^2 .

So, the resulting potential is seen to be energy-dependent. We make our modification to M and then K so that only an energy-independent potential arises which is more in keeping with our usual intuition for understanding the occurrence of bound states.

As the k^0 -dependent part of the effective potential is associated with a temporal distance-one κ^6 contribution to K , we modify M so that this contribution is smaller. This happens because M is modified so that $M^{(6)}$ has the same product structure as $M_0^{(6)}$. By doing so, the decay of the modified K is improved for temporal distance one. In contrast to the unmodified M , for our modified $M \equiv M'$, we get a $\kappa^8 |x_3^0 - x_1^0|$ decay for any distance $|x_3^0 - x_1^0| \geq 1$. This comes about as the modified $M^{(6)}$ also has a product structure so that the modified $K(x_1, x_2, x_3, x_4)$ in Eq. (4.28) has a zero sixth derivative $K^{(6)} = 0$, for $x_1^0 = x_2^0 < x_3^0 = x_4^0$, since the last two terms on the r.h.s. of Eq. (D5) have now the same coefficient as the first term.

There is still a possibility of having a κ^8 contribution to the modified $K(x_1, x_2, x_3, x_4)$ of temporal distance one leading to an energy-dependent potential of order κ^2 , which would be of the same order of the quasi-meson exchange potential. These κ^8 contributions arise from M and M_0 with temporal distance one, but vanish due to imbalance of fermion components.

With this modification, the space-range zero potential is zero and the dominant contribution is the quasimeson space-range one exchange potential.

APPENDIX E: Determination of the space range-one exchange potential

In this Appendix, we determine the quasi-meson exchange potential 20×20 matrix of Section 4.3 in the individual spin basis, i.e. $v_{s_1 s_2 s_3 s_4}^{(2)}$, $v_{s_1 s_2 t_3 t_4}^{(2)}$, $v_{t_1 t_2 s_3 s_4}^{(2)}$, and $v_{t_1 t_2 t_3 t_4}^{(2)}$. We give reduced formulas which we use for computation, and present the derivation explicitly for $v_{s_1 s_2 s_3 s_4}^{(2)}$ given in Eq. (4.31), the elements of which, in the total spin basis, are given below Eq. (4.32). The other elements are obtained in an analogous manner.

In Eq. (4.31), consider the $\kappa = 0$ averages expressed as a sum of all contractions using Wick's theorem. If the explicit $\bar{\psi}\psi$ contract in the first average then $\alpha_1 = \beta_2$ and $\Gamma_{\alpha_1 \beta_1}^{e^1} \Gamma_{\alpha_2 \alpha_1}^{-e^1} = 0$; the same for the second average. Thus, only contractions involving one explicit internal $\bar{\psi}$ and one quark field of $\bar{B}$'s contribute. Also, recalling that the ψ 's on the $\bar{B}$'s only have lower spin indices, this forces the spin index of the explicit $\bar{\psi}$'s to be also lower in order not to get a zero contribution. Next, noting that in the subspace of lower components the spin matrix $\Gamma^{\pm e^1}$ reduces to minus the identity, and after using the $\kappa = 0$

isospin flip symmetry in the second factor, we can write

$$v_{s_1, s_2, s_3, s_4}^{(2)} = \sum_{I_1, I_3} c_{I_1 - I_1}^{\frac{1}{2} \frac{1}{2}} c_{I_3 - I_3}^{\frac{1}{2} \frac{1}{2}} \times \langle B_{\frac{1}{2} I_1 s_1} \bar{B}_{\frac{1}{2} I_3 s_3} : \bar{\psi}_{a\alpha f_1} \psi_{a\beta f_2} : \rangle^{(0)} \times \langle B_{\frac{1}{2} I_1 s_2} \bar{B}_{\frac{1}{2} I_3 s_4} : \bar{\psi}_{b\beta - f_2} \psi_{b\alpha - f_1} : \rangle^{(0)}, \quad (\text{E1})$$

with all fields at a same site and where the Wick order $::$ forbids contraction between the explicit internal $\bar{\psi}$'s. To implement the exclusion of this contraction, we note that up to a global sign and omitting the detailed field indices and summations we have $\langle B \bar{B} \psi \bar{\psi} \rangle^{(0)} \sim \langle \psi_1 \psi_2 \psi_3 \bar{\psi}_5 \bar{\psi}_6 \bar{\psi}_7 \psi_4 \bar{\psi}_8 \rangle^{(0)} \sim \langle \psi_1 \psi_2 \psi_3 \psi_4 \bar{\psi}_5 \bar{\psi}_6 \bar{\psi}_7 \bar{\psi}_8 \rangle^{(0)} = \det[\delta_{ij}]$, $i = 1, \dots, 4$ and $j = 5, \dots, 8$, such that the Wick order in $:\psi_4 \bar{\psi}_8:$ is equivalent to replacing δ_{78} by zero in the previous fermionic Gram determinant $\det[\delta_{ij}]$.

To continue, we use symmetries and isospin and spin restrictions to reduce the number of $v^{(2)}$'s to be calculated. Fixing a matrix element in $v^{(2)}$, we consider the terms in the sums labeled by $(I_1 I_3)$, $(I_1 J_3)$, $(J_1 I_3)$ or $(J_1 J_3)$. Isospin flip symmetry gives equality of the $(J_1 J_3)$ and $(-J_1 - J_3)$ terms, etc..., where we take into account the properties of the C-G coefficients. Isospin conservation forces the restrictions $I_1 + f_2 = I_3 + f_1 \Rightarrow |I_1 - I_3| \leq 1$, $I_1 + f_2 = J_3 + f_1 \Rightarrow |I_1 - J_3| \leq 1$, $J_1 + f_2 = J_3 + f_1 \Rightarrow |J_1 - J_3| \leq 1$. In addition, we have spin sum restrictions $s_1 - s_3 = s_4 - s_2 = \alpha - \beta$, $|s_1 - s_3| \leq 1$; $s_1 - t_3 = t_4 - s_2 = \alpha - \beta$, $|s_1 - t_3| \leq 1$; $t_1 - t_3 = t_4 - t_2 = \alpha - \beta$, $|t_1 - t_3| \leq 1$. Also, for $\kappa = 0$, conjugation (denoted by $\mathcal{C}_0$) is a symmetry. Using $\mathcal{C}_0$ gives the equality of the $(I_1 I_3)$ and $(I_3 I_1)$ ($(J_1 J_3)$ and $(J_3 J_1)$) product of summed expectations if $s_1 = s_3$ and $s_2 = s_4$ ($t_1 = t_3$ and $t_2 = t_4$), such as diagonal elements. In this way, we show that many of the terms are zero. We can also, after applying $\mathcal{C}_0$ to both factors, change $f_1 \leftrightarrow -f_1$, $f_2 \leftrightarrow -f_2$ to get equality of the $(I_1 I_3)$ and $(I_3 I_1)$ terms if $s_1 = s_4$ and $s_2 = s_3$; similarly for the J 's. We can also impose the prohibited contraction condition to show that certain elements are zero.

In addition to the fact that the 20×20 potential matrix is symmetric there are other symmetries. By making the change of labeling $\alpha \leftrightarrow \beta$, $f_1 \leftrightarrow -f_2$, $f_2 \leftrightarrow -f_1$, the following symmetry properties are obtained

$$v_{s_1 s_2 s_3 s_4}^{(2)} = v_{s_2 s_1 s_4 s_3}^{(2)}, \quad v_{s_1 s_2 t_3 t_4}^{(2)} = v_{s_2 s_1 t_4 t_3}^{(2)}, \\ v_{t_1 t_2 t_3 t_4}^{(2)} = v_{t_2 t_1 t_4 t_3}^{(2)}.$$

Also, the spin flip change $\alpha \leftrightarrow -\alpha$, $\beta \leftrightarrow -\beta$ gives the equality of the $v^{(2)}$'s for $s_k \rightarrow -s_k$, $t_j \rightarrow -t_j$, for all j 's and k 's.

Thus, there is symmetry in the upper left 4×4 (lower right 16×16) matrix about the secondary diagonal of the submatrix.

Using each symmetry alone or coupling them explains the repeated values occurring in the matrix with the following two exceptions: the equality of the (69) and (811) elements and the equality between (1111) and

(1114). By spin flip at $\kappa = 0$, we see that (11) = (44), (22) = (33), (17) = (418), (110) = (415), (113) = (412), (28) = (317), (211) = (314), (214) = (311); by spin flip at $\kappa = 0$ and $1 \leftrightarrow 2$, $3 \leftrightarrow 4$ exchanges (113) = (418), (77) = (1212); by $1 \leftrightarrow 2$, $3 \leftrightarrow 4$ exchanges (66) = (99). By applying $\mathcal{C}_0$ to both average factors, we obtain (710) = (1013). Thus, we only have to calculate e.g. the elements (11), (22), (23), (17), (110), (113), (28), (211), (214), (55), (66), (69), (77), (710), (88), (811), (1010), (1111) and (1114).

The values of the above elements are given below Eq. (4.32). As a check in calculating the elements of Eq. (4.31), etc, it is useful to give the decomposition of the total value before performing the sums in the equivalent expression of Eq. (E1). We now give the nonzero values for the partial terms $(I_1 I_3)$ or $(I_1 J_3)$ or $(J_1 J_3)$ for each matrix element; the values for $(-I_1 - I_3)$ or $(-I_1 - J_3)$ or $(-J_1 - J_3)$ are the same. The sum of the values for each relevant partial matrix element given below multiplied by a factor 2 is the value of the matrix element. The values, given here with precision of two digits, are as follows. For the matrix element (11), which corresponds to $s_1 = s_2 = 1/2$, $s_3 = s_4 = 1/2$, in Eq. (E1), we need the contributions $(I_1 I_3) = (\frac{1}{2} \frac{1}{2})$, which is 0.78, and $(I_1 I_3) = (\frac{1}{2} - \frac{1}{2})$, which is -0.94. Denoting this contribution as (11): $(\frac{1}{2} \frac{1}{2})$ 0.78, $(\frac{1}{2} - \frac{1}{2})$ -0.94, and using this notation for the other elements we find (22): $(\frac{1}{2} \frac{1}{2})$ 1.22, $(\frac{1}{2} - \frac{1}{2})$ 0.44; (23): $(\frac{1}{2} \frac{1}{2})$ -0.44, $(\frac{1}{2} - \frac{1}{2})$ -1.39; (17): $(\frac{1}{2} \frac{3}{2})$ -0.41, $(\frac{1}{2} \frac{1}{2})$ -0.27, $(\frac{1}{2} - \frac{1}{2})$ -0.14; (110): $(\frac{1}{2} \frac{3}{2})$ 0.47, $(\frac{1}{2} \frac{1}{2})$ 0.31, $(\frac{1}{2} - \frac{1}{2})$ 0.16; (113): $(\frac{1}{2} \frac{3}{2})$ -0.41, $(\frac{1}{2} \frac{1}{2})$ -0.27, $(\frac{1}{2} - \frac{1}{2})$ -0.14; (28): $(\frac{1}{2} \frac{3}{2})$ 0.71, $(\frac{1}{2} \frac{1}{2})$ 0.47, $(\frac{1}{2} - \frac{1}{2})$ 0.24; (211): $(\frac{1}{2} \frac{3}{2})$ -0.47, $(\frac{1}{2} \frac{1}{2})$ -0.31, $(\frac{1}{2} - \frac{1}{2})$ -0.16; (214): $(\frac{1}{2} \frac{3}{2})$ 0.24, $(\frac{1}{2} \frac{1}{2})$ 0.16, $(\frac{1}{2} - \frac{1}{2})$ 0.08; (55): $(\frac{3}{2} \frac{1}{2})$ -0.75, $(\frac{1}{2} \frac{3}{2})$ -0.75, $(\frac{1}{2} \frac{1}{2})$ 1.00, $(\frac{1}{2} - \frac{1}{2})$ -1.00; (66): $(\frac{3}{2} \frac{1}{2})$ -0.50, $(\frac{1}{2} \frac{3}{2})$ -0.50, $(\frac{1}{2} \frac{1}{2})$ 0.67, $(\frac{1}{2} - \frac{1}{2})$ -0.67; (69): $(\frac{3}{2} \frac{1}{2})$ -0.25, $(\frac{1}{2} \frac{3}{2})$ -0.25, $(\frac{1}{2} \frac{1}{2})$ 0.33, $(\frac{1}{2} - \frac{1}{2})$ -0.33; (77): $(\frac{3}{2} \frac{1}{2})$ -0.25, $(\frac{1}{2} \frac{3}{2})$ -0.25, $(\frac{1}{2} \frac{1}{2})$ 0.33, $(\frac{1}{2} - \frac{1}{2})$ -0.33; (710): $(\frac{3}{2} \frac{1}{2})$ -0.29, $(\frac{1}{2} \frac{3}{2})$ -0.29, $(\frac{1}{2} \frac{1}{2})$ 0.38, $(\frac{1}{2} - \frac{1}{2})$ -0.38; (88): all zero; (811): $(\frac{3}{2} \frac{1}{2})$ -0.25, $(\frac{1}{2} \frac{3}{2})$ -0.25, $(\frac{1}{2} \frac{1}{2})$ 0.33, $(\frac{1}{2} - \frac{1}{2})$ -0.33; (1010): $(\frac{3}{2} \frac{1}{2})$ -0.42, $(\frac{1}{2} \frac{3}{2})$ -0.42, $(\frac{1}{2} \frac{1}{2})$ 0.56, $(\frac{1}{2} - \frac{1}{2})$ -0.56; (1111): $(\frac{3}{2} \frac{1}{2})$ -0.33, $(\frac{1}{2} \frac{3}{2})$ -0.33, $(\frac{1}{2} \frac{1}{2})$ 0.44, $(\frac{1}{2} - \frac{1}{2})$ -0.44; (1114): $(\frac{3}{2} \frac{1}{2})$ -0.33, $(\frac{1}{2} \frac{3}{2})$ -0.33, $(\frac{1}{2} \frac{1}{2})$ 0.44, $(\frac{1}{2} - \frac{1}{2})$ -0.44.

The values given below Eq. (4.32) are the exact values for $v^{(2)}$ obtaining by summing up the exact values for the above partial decomposition values and multiplying by two.

APPENDIX F: Uncorrelated four-baryon function and spectral representation

Here we show how to obtain a spectral representation for $\hat{M}_{0, \ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3, x_4)$, and its total spin version; the components of which are obtained by summing over the appropriate C-G combinations of the $\hat{F}_{0, \ell_1 \ell_2 \ell_3 \ell_4}(x_1, x_2, x_3, x_4)$. Our starting point is Eq.

(4.24):

$$F_{0,1234} = -G_{13}G_{24} + G_{14}G_{23},$$

where we use a shorthand notation for the collective indices (ℓ, x) .

In lattice relative coordinates and in the equal time representation

$$F_{0,1234}(\vec{\xi}, \vec{\eta}, \tau) = -G_{13}(\tau + \vec{\xi}) G_{24}(\tau + \vec{\eta}) + G_{14}(\tau + \vec{\xi} + \vec{\eta}) G_{23}(\tau), \quad (\text{F1})$$

and setting $\vec{k} = \vec{0}$ in the Fourier transform of τ only, with dual variable k , our goal is to obtain a spectral representation for

$$\hat{F}_{0,1234}(\vec{\xi}, \vec{\eta}, k^0) = \int F_{0,1234}(\vec{\xi}, \vec{\eta}, \tau) e^{-ik^0 \tau} d\tau.$$

For $x^0 \neq 0$, and using the spectral representation for $G_{ij}(x)$ of Eq. (3.11), we obtain

$$G_{ij}(x) = - \int_{-1}^1 \int_{\mathbb{T}^3} (\lambda^0)^{|x^0|-1} e^{i\vec{p} \cdot \vec{x}} d_{\lambda^0} \alpha_{\vec{p},ij}(\lambda^0) d\vec{p}, \quad (\text{F2})$$

where, by parity, $\alpha_{\vec{p},ij} = \alpha_{-\vec{p},ij}$.

Separating the $\tau^0 = 0$ contribution and substituting the spectral representation for the G 's into Eq. (F1), after performing the integral over τ , we get

$$\begin{aligned} \hat{F}_{0,1234}(\vec{\xi}, \vec{\eta}, k^0) = & (2\pi)^{-3} \int_{\mathbb{T}^3} \left[-\tilde{G}_{13}(\vec{p}) \tilde{G}_{24}(\vec{p}) \Xi_+(\vec{p}) \right. \\ & \left. + \tilde{G}_{14}(\vec{p}) \tilde{G}_{23}(\vec{p}) \Xi_-(\vec{p}) \right] d\vec{p} \\ & + (2\pi)^3 \int_{-1}^1 \int_{-1}^1 \int_{\mathbb{T}^3} f(k^0, \lambda^0 \lambda'^0) \\ & \times \left[-d_{\lambda^0} \alpha_{\vec{p},13}(\lambda^0) d_{\lambda'^0} \alpha_{\vec{p},24}(\lambda'^0) \Xi_+(\vec{p}) \right. \\ & \left. + d_{\lambda^0} \alpha_{\vec{p},14}(\lambda^0) d_{\lambda'^0} \alpha_{\vec{p},23}(\lambda'^0) \Xi_-(\vec{p}) \right] d\vec{p}, \quad (\text{F3}) \end{aligned}$$

where we recall that $\tilde{G}(\vec{p}) = \sum_{\vec{x}} e^{-i\vec{p} \cdot \vec{x}} G(x^0 = 0, \vec{x})$, and where we have

$$\Xi_{\pm}(\vec{p}) \equiv \Xi_{\pm}(\vec{p}, \vec{\xi}, \vec{\eta}) = \cos \vec{p} \cdot \vec{\xi} \cos \vec{p} \cdot \vec{\eta} \pm \sin \vec{p} \cdot \vec{\xi} \sin \vec{p} \cdot \vec{\eta}.$$

To proceed in our deduction of $\hat{M}_0$, we make the approximation that $\tilde{G}$ is diagonal in spin, and independent of the isospin I and spin (which holds up to and including $\mathcal{O}(\kappa^5)$, see Ref. [5]), and we denote it simply by $\tilde{G}(\vec{p})$, with no index; and the same for the measure $d_{\lambda^0} \alpha_{\vec{p}}$.

Under this hypothesis, carrying out the sum over the isospin C-G coefficients we obtain, for the 11 element of $\hat{M}_0$ in the individual spin basis

$$\begin{aligned} \hat{M}_{0,11,s_1 s_2 s_3 s_4}(\vec{\xi}, \vec{\eta}, k^0) = & (2\pi)^{-3} \int_{\mathbb{T}^3} [\tilde{G}(\vec{p})]^2 [-\delta_{s_1 s_3} \delta_{s_2 s_4} \Xi_+(\vec{p}) \\ & - \delta_{s_1 s_4} \delta_{s_2 s_3} \Xi_-(\vec{p})] d\vec{p} \\ & + (2\pi)^3 \int_{-1}^1 \int_{-1}^1 \int_{\mathbb{T}^3} f(k^0, \lambda^0 \lambda'^0) \times \\ & [-\delta_{s_1 s_3} \delta_{s_2 s_4} \Xi_+(\vec{p}) - \delta_{s_1 s_4} \delta_{s_2 s_3} \Xi_-(\vec{p})] \\ & \times d_{\lambda^0} \alpha_{\vec{p}}(\lambda^0) d_{\lambda'^0} \alpha_{\vec{p}}(\lambda'^0) d\vec{p}. \quad (\text{F4}) \end{aligned}$$

The 22 element of $\hat{M}_0$ is obtained in the same way, by replacing s_j by t_j in the above. The off-diagonal elements are zero.

Finally, we obtain the total spin basis representation for $\hat{M}_0$ by carrying out the sums over the spin C-G coefficients and using the explicit values and symmetry properties of the C-G coefficients, we obtain the expression given in Eq. (4.36).

-
- [1] K. Wilson, in *New Phenomena in Subnuclear Physics*, Part A, A. Zichichi ed. (Plenum Press, NY, 1977).
 - [2] M. Creutz, *Quarks, Gluons and Lattices* (Cambridge University Press, Cambridge, 1983).
 - [3] I. Montvay and G. Münster, *Quantum Fields on a Lattice* (Cambridge University Press, Cambridge, 1997).
 - [4] P.A. Faria da Veiga, M. O'Carroll, and R. Schor, Phys. Rev. **D 67**, 017501 (2003).
 - [5] P.A. Faria da Veiga, M. O'Carroll, and R. Schor, Commun. Math. Phys. **245**, 383 (2004).
 - [6] A. Francisco Neto, P.A. Faria da Veiga, and M. O'Carroll, J. Math. Phys. **45**, 628 (2004).
 - [7] P.A. Faria da Veiga, M. O'Carroll, and R. Schor, Phys. Rev. **D 68**, 037501 (2003).
 - [8] P.A. Faria da Veiga, M. O'Carroll, and A. Francisco Neto, Phys. Rev. **D 69**, 097501 (2004).
 - [9] P.A. Faria da Veiga and M. O'Carroll, Phys. Rev. **D 71**, 017503 (2005).
 - [10] P.A. Faria da Veiga, M. O'Carroll, and A. Francisco Neto, Phys. Rev. **D 72**, 034507 (2005).
 - [11] T. Banks et al., Phys. Rev. **D 15**, 1111 (1977).
 - [12] K. Osterwalder and E. Seiler, Ann. Phys. (NY) **110**, 440 (1978).
 - [13] E. Seiler, Lect. Notes in Phys. **159**, *Gauge Theories as a Problem of Constructive Quantum Field Theory and Statistical Mechanics* (Springer, New York, 1982).
 - [14] J. Fröhlich and C. King, Nucl. Phys **B 290**, 157 (1987).
 - [15] I. Montvay, Rev. Mod. Phys. **59**, 263 (1987).
 - [16] F. Myhrer and J. Wroldsen, Rev. Mod. Phys. **60**, 629 (1988).
 - [17] D. Schreiber, Phys. Rev. **D 48**, 5393 (1993).
 - [18] M. Creutz, Nucl. Phys. **B** (Proc. Suppl.) **94** 219 (2001).
 - [19] R. Machleidt, Nucl. Phys. **A 689**, 11c (2001).
 - [20] R. Machleidt, K. Holinde, and Ch. Elster, Phys. Rep.

- 149**, 1 (1987).
- [21] M.J. Savage, *Nuclei from QCD : Strategy, challenges and status*, PANIC 05, arXiv:nucl-th/0601001.
 - [22] H.R. Fiebig, H. Markum, A. Mihály, and K. Rabitsch, Nucl. Phys. Proc. Suppl. **53**, 804 (1997).
 - [23] C. Stewart and R. Koniuk, Phys. Rev. **D 57**, 5581 (1998).
 - [24] H.R. Fiebig and H. Markum in *International Review of Nuclear Physics, Hadronic Physics from Lattice QCD*, A.M. Green ed. (World Scientific, Singapoure, 2003).
 - [25] Ph. de Forcrand and S. Kim, *The Spectrum of lattice QCD with staggered fermions at strong coupling* (2006), hep-lat/0608012.
 - [26] R.S. Schor and M. O'Carroll, Phys. Rev. **E 62**, 1521 (2000). J. Stat. Phys. **99**, 1207 (2000); **99**, 1265 (2000); and **109**, 279 (2002).
 - [27] P.A. Faria da Veiga, M. O'Carroll, E. Pereira, and R. Schor, Commun. Math. Phys. **220**, 377(2001).
 - [28] M. Reed and B. Simon, *Modern Methods of Mathematical Physics*, vol. 4, *Analysis of Operators*, (Academic Press, New York, 1978).
 - [29] P.A. Faria da Veiga and M. O'Carroll, (2006).
 - [30] T. Spencer, Commun. Math. Phys. **44**, 143 (1975).
 - [31] B. Simon, *Representations of Finite and Compact Groups* (American Mathematical Society, Providence, 1996).
 - [32] F.A. Berezin, *The Method of Second Quantization* (Academic Press, NY, 1966).
 - [33] B. Simon, *Statistical Mechanics of Lattice Models* (Princeton University Press, Princeton, 1994).
 - [34] P.H.R. dos Anjos, P.A. Faria da Veiga, and M. O'Carroll, in preparation.
 - [35] J. Dimock and J.-P. Eckmann, Commun. Math. Phys. **51**, 41 (1976); Annals of Physics (NY) **103**, 289 (1977).
 - [36] M. Reed and B. Simon, *Modern Methods of Mathematical Physics*, vol. 2, *Fourier Analysis, Self-Adjointness* (Academic Press, New York, 1975).
 - [37] A.A. Maradudin, E.W. Montroll, G.H. Weiss, *Theory of Lattice Dynamics in the Harmonic Approximation*, in *Solid State Physics*, supplement 3, F. Seitz and D. Turnbull eds., (Academic Press, New York, 1963).
 - [38] E.M. Lifchitz, Nouvo Cimento Suppl. **10 3**, 716 (1956).
 - [39] W.A. Bowers and H.B. Rosenstock, J. Chem. Phys. **18**, 1056 (1950).
 - [40] G.F. Koster, Phys. Rev. **95**, 1436 (1954).
 - [41] S. Katsura, S. Inawashiro, J. Math. Phys. **12**, 1622 (1971).
 - [42] R.T. Delves, G.S. Joyce, Ann. Phys. (NY) **291**, 71 (2001).
 - [43] P.A. Faria da Veiga, L. Ioriatti, and M. O'Carroll, Phys. Rev. E **66**, 016130 (2002).
 - [44] P.A. Faria da Veiga, and M. O'Carroll, in JHEP Proceedings, PrHEP UNESP2002, *Workshop on integrable theories, solitons and duality*, L.A. Ferreira ed., São Paulo (2002).
 - [45] J. Glimm and A. Jaffe, *Quantum Physics: A Functional Integral Point of View* (Springer Verlag, New York, 1986).
 - [46] Precisely, the complexification of κ_p is done by a replacement of a finite number of parameters in the polymer activities of the polymer expansion representation of the correlation functions. Note that, with this, we take care of the normalizations of correlations, and the complex conjugation operation present in averages $\langle \cdot \rangle$ are taken before complexification of κ_p so that only dependence in $\kappa_p \in \mathbb{C}$ occurs (*not* in κ_p^* !).
 - [47] E. Hille, *Analytic Function Theory*, vol. 1 (Blaisdell Publishing Co., New York, 1959).
 - [48] R.S. Schor and M. O'Carroll, Commun. Math. Phys. **103**, p.569 (1986).